

REALIZATION OF METRIC SPACES AS INVERSE LIMITS, AND BILIPSCHITZ EMBEDDING IN L_1

JEFF CHEEGER AND BRUCE KLEINER

ABSTRACT. We give sufficient conditions for a metric space to bilipschitz embed in L_1 . In particular, if X is a length space and there is a Lipschitz map $u : X \rightarrow \mathbb{R}$ such that for every interval $I \subset \mathbb{R}$, the connected components of $u^{-1}(I)$ have diameter $\leq \text{const} \cdot \text{diam}(I)$, then X admits a bilipschitz embedding in L_1 . As a corollary, the Laakso examples [Laa00] bilipschitz embed in L_1 , though they do not embed in any Banach space with the Radon-Nikodym property (e.g. the space ℓ_1 of summable sequences).

The spaces appearing the statement of the bilipschitz embedding theorem have an alternate characterization as inverse limits of systems of metric graphs satisfying certain additional conditions. This representation, which may be of independent interest, is the initial part of the proof of the bilipschitz embedding theorem. The rest of the proof uses the combinatorial structure of the inverse system of graphs and a diffusion construction, to produce the embedding in L_1 .

CONTENTS

1. Introduction	2
2. Notation and preliminaries	12
3. Inverse systems of graphs with path metrics	18
4. Realizing metric spaces as limits of admissible inverse systems	20
5. A special case of Theorem 1.16	22
6. Slices and the associated measures	22
7. Estimates on the family of measures $\{\Sigma'_i\}_{i \in \mathbb{Z}}$	28

Date: October 12, 2011.

J.C. supported by NSF grants DMS-0704404 and DMS-1005552. B.K. supported by NSF grants DMS-1007508 and DMS-1105656.

8. Proof of Theorem 1.16 under Assumption 5.1	34
9. The proof of Theorem 1.16, general case	35
10. The Laakso examples from [Laa00] and Example 1.4	39
11. Realizing metric spaces as inverse limits: further generalization	41
References	43

1. INTRODUCTION

Overview. This paper is part of a series [CK06b, CK06a, CK10a, CK09, CK10b, CKN09, CKc] which examines the relations between differentiability properties and bilipschitz embeddability in Banach spaces. We give a new criterion for metric spaces to bilipschitz embed in L_1 . This applies to several known families of spaces, illustrating the sharpness of earlier nonembedding theorems. In the first part of the proof, we characterize a certain class of metric spaces as inverse limits; this may be of independent interest.

Metric spaces sitting over \mathbb{R} . We begin with a special case of our main embedding theorem.

Theorem 1.1. *Let X be a length space. Suppose $u : X \rightarrow \mathbb{R}$ is a Lipschitz map, and there is a $C \in (0, \infty)$ such that for every interval $I \subset \mathbb{R}$, each connected component of $u^{-1}(I)$ has diameter at most $C \cdot \text{diam}(I)$. Then X admits a bilipschitz embedding $f : X \rightarrow L_1(Z, \mu)$, for some measure space (Z, μ) .*

We illustrate Theorem 1.1 with two simple examples:

Example 1.2 (Lang-Plaut [LP01], cf. Laakso [Laa00]). We construct a sequence of graphs $\{X_i\}_{i \geq 0}$ where X_i has a path metric so that every edge has length 4^{-i} . Let X_0 be the unit interval $[0, 1]$. For $i > 0$, inductively construct a X_i from X_{i-1} by replacing each edge of X_{i-1} with a copy of the graph Γ in Figure 1, rescaled by the factor $m^{-(i-1)}$. The graphs X_1 , X_2 , and X_3 are shown. The sequence $\{X_i\}$ naturally forms an inverse system,

$$X_0 \xleftarrow{\pi_0} \cdots \xleftarrow{\pi_i} X_i \xleftarrow{\pi_{i+1}} \cdots,$$

where the projection map $\pi_{i-1} : X_i \rightarrow X_{i-1}$ collapses the copies of Γ to intervals. The inverse limit X_∞ has a metric d_∞ given by

$$(1.3) \quad d_\infty(x, x') = \lim_{i \rightarrow \infty} d_{X_i}(\pi_i^\infty(x), \pi_i^\infty(x')),$$

where $\pi_i^\infty : X_\infty \rightarrow X_i$ denotes the canonical projection. (Note that the sequence of metric spaces $\{X_i\}_{i \geq 0}$ Gromov-Hausdorff converges to (X_∞, d_∞) .) It is not hard to verify that $\pi_0^\infty : (X_\infty, d_\infty) \rightarrow [0, 1]$ satisfies the hypotheses of Theorem 1.1.




FIGURE 1.

Example 1.4. Construct an inverse system

$$X_0 \xleftarrow{\pi_0} \dots \xleftarrow{\pi_i} X_i \xleftarrow{\pi_{i+1}} \dots$$

inductively as follows. Let $X_0 = [0, 1]$. For $i > 0$, inductively define X'_{i-1} to be the result of trisecting all edges in X_{i-1} , and let $N \subset X'_{i-1}$ be new vertices added in trisection. Now form X_i by taking two copies of X'_{i-1} and gluing them together along N . More formally,

$$X_i = (X'_{i-1} \times \{0, 1\}) / \sim,$$

where $(v, 0) \sim (v, 1)$ for all $v \in N$. The map $\pi_{i-1} : X_i \rightarrow X_{i-1}$ is induced by the collapsing map $X'_{i-1} \times \{0, 1\} \ni (x, j) \mapsto x \in X_{i-1}$. Metrizing the inverse limit X_∞ as in Example 1.2, the canonical projection $X_\infty \rightarrow X_0 \simeq [0, 1] \rightarrow \mathbb{R}$ satisfies the assumptions of Theorem 1.1.

The inverse limit X_∞ in Example 1.4 is actually bilipschitz homeomorphic to one of the Ahlfors regular Laakso spaces from [Laa00], see Section 10. Thus Theorem 1.1 implies that this Laakso space bilipschitz embeds in L_1 (this special case was announced in [CK06b]). Laakso showed that X_∞ carries a doubling measure which satisfies a Poincaré inequality, and using this, the nonembedding result of [CK09] implies

that X_∞ does not bilipschitz embed in any Banach space which satisfies the Radon-Nikodym property. Therefore we have:

Corollary 1.5. *There is a compact Ahlfors regular (in particular doubling) metric measure space satisfying a Poincaré inequality, which bilipschitz embeds in L_1 , but not in any Banach space with the Radon-Nikodym property (such as ℓ_1).*

To our knowledge, this is the first example of a doubling space which bilipschitz embeds in L_1 but not in ℓ_1 .

We can extend Theorem 1.1 by dropping the length space condition, and replacing connected components with a metrically based variant.

Definition 1.6. Let Z be a metric space and $\delta \in (0, \infty)$. A δ -**path** (or δ -**chain**) in Z is a finite sequence of points $z_0, \dots, z_k \subset Z$ such that $d(z_{i-1}, z_i) \leq \delta$ for all $i \in \{1, \dots, k\}$. The property of belonging to a δ -path defines an equivalence relation on Z , whose cosets are the δ -**components** of Z .

Our main embedding result is:

Theorem 1.7. *Let X be a metric space. Suppose there is a 1-Lipschitz map $u : X \rightarrow \mathbb{R}$ and a constant $C \in (0, \infty)$ such that for every interval $I \subset \mathbb{R}$, the $\text{diam}(I)$ -components of $u^{-1}(I)$ have diameter at most $C \cdot \text{diam}(I)$. Then X admits a bilipschitz embedding $f : X \rightarrow L_1$.*

Inverse systems of directed metric graphs, and multi-scale factorization. Our approach to proving Theorem 1.7 is to first show that any map $u : X \rightarrow \mathbb{R}$ satisfying the hypothesis of theorem can be factored into an infinite sequence of maps, i.e. it gives rise to a certain kind of inverse system where X reappears (up to bilipschitz equivalence) as the inverse limit. Strictly speaking this result has nothing to do with embedding, and can be viewed as a kind of multi-scale version of monotone-light factorization ([Eil34, Why34]) in the metric space category.

We work with a special class of inverse systems of graphs:

Definition 1.8 (Admissible inverse systems). An inverse system indexed by the integers

$$\dots \xleftarrow{\pi_{-i-1}} X_{-i} \xleftarrow{\pi_{-i}} \dots \xleftarrow{\pi_{-1}} X_0 \xleftarrow{\pi_0} \dots \xleftarrow{\pi_i} X_i \xleftarrow{\pi_{i+1}} \dots,$$

is **admissible** if for some integer $m \geq 2$ the following conditions hold:

- (1) X_i is a nonempty directed graph for every $i \in \mathbb{Z}$.
- (2) For every $i \in \mathbb{Z}$, if X'_i denotes the directed graph obtained by subdividing each edge of X_i into m edges, then π_i induces a map $\pi_i : X_{i+1} \rightarrow X'_i$ which is simplicial, an isomorphism on every edge, and direction preserving.
- (3) For every $i, j \in \mathbb{Z}$, and every $x \in X_i$, $x' \in X_j$, there is a $k \leq \min(i, j)$ such that x and x' project to the same connected component of X_k .

Note that the X_i 's need not be connected or have finite valence, and they may contain isolated vertices.

We endow each X_i with a (generalized) path metric $d_i : X_i \times X_i \rightarrow [0, \infty]$, where each edge is linearly isometric to the interval $[0, m^{-i}] \subset \mathbb{R}$. Since we do not require the X_i 's to be connected, we have $d_i(x, x') = \infty$ when x, x' lie in different connected components of X_i . It follows from Definition 1.8 that the projection maps $\pi_i^j : (X_j, d_j) \rightarrow (X_i, d_i)$ are 1-Lipschitz.

Examples 1.2 and 1.4 provide admissible inverse systems in a straightforward way: for $i < 0$ one simply takes X_i to be a copy of \mathbb{R} with the standard subdivision into intervals of length m^{-i} , and the projection map $\pi_i : X_{i+1} \rightarrow X_i$ to be the identity map. Of course this modification does not affect the inverse limit.

Let X_∞ be the inverse limit of the system $\{X_i\}$, and let $\pi_i^j : X_j \rightarrow X_i$, $\pi_i^\infty : X_\infty \rightarrow X_i$ denote the canonical projections for $i \leq j \in \mathbb{Z}$. We will often omit the superscripts and subscripts when there is no risk of confusion.

We now equip the inverse limit X_∞ with a metric \bar{d}_∞ ; unlike in the earlier examples, this is not defined as a limit of pseudo-metrics $d_i \circ \pi_i^\infty$.

Definition 1.9. Let $\bar{d}_\infty : X_\infty \times X_\infty \rightarrow [0, \infty)$ be the supremal pseudo-distance on X_∞ such that for every $i \in \mathbb{Z}$ and every vertex $v \in X_i$, if

$$\text{St}(v, X_i) = \cup\{e \mid e \text{ is an edge of } X_i, v \in e\}$$

is the closed star of v in X_i , then the inverse image of $\text{St}(v, X_i)$ under the projection map $X_\infty \rightarrow X_i$ has diameter at most $2m^{-i}$. Henceforth, unless otherwise indicated, distances in X_∞ will refer to \bar{d}_∞ .

In fact \bar{d}_∞ is a metric, and for any distinct points $x, x' \in X_\infty$, the distance $\bar{d}_\infty(x, x')$ is comparable to m^{-i} , where i is the maximal integer

such that $\{\pi_i^\infty(x), \pi_i^\infty(x')\}$ is contained in the star of some vertex $v \in X_i$; see Section 2. In Examples 1.2 and 1.4, the metric \bar{d}_∞ is comparable to the metric d_∞ defined using the path metrics in (1.3); see Section 3.

Admissible inverse systems give rise to spaces satisfying the hypotheses of Theorem 1.7:

Theorem 1.10. *Let $\{X_i\}$ be an admissible inverse system. Then there is a 1-Lipschitz map $\phi : X_\infty \rightarrow \mathbb{R}$ which is canonical up to post-composition with a translation, which satisfies the assumptions of Theorem 1.7.*

The converse is also true:

Theorem 1.11. *Let X be a metric space. Suppose $u : X \rightarrow \mathbb{R}$ is a 1-Lipschitz map, and there is a constant $C \in [1, \infty)$ such that for every interval $I \subset \mathbb{R}$, the inverse image $u^{-1}(I) \subset X$ has $\text{diam}(I)$ -components of diameter at most $C \cdot \text{diam}(I)$. Then for any $m \geq 2$ there is an admissible inverse system $\{X_i\}$ and a compatible system of maps $f_i : X \rightarrow X_i$, such that:*

- *The induced map $f_\infty : X \rightarrow (X_\infty, \bar{d}_\infty)$ is $L' = L'(C, m)$ -bilipschitz.*
- *$u = \phi \circ f_\infty$, where $\phi : X_\infty \rightarrow \mathbb{R}$ is the 1-Lipschitz map of Theorem 1.10.*

Theorem 1.1 is a corollary of Theorem 1.11 : if $u : X \rightarrow \mathbb{R}$ is as in Theorem 1.1, then for any interval $[a, a + r] \subset \mathbb{R}$, an r -component of $f^{-1}([a, a + r])$ will be contained in a connected component of $f^{-1}([a - r, a + 2r])$ (since X is a length space), and therefore has diameter $\leq 3C \text{diam}(I)$.

Remark 1.12. Theorem 1.11 implies that Examples 1.2 and 1.4 can be represented up to bilipschitz homeomorphism as inverse limits of many different admissible inverse systems, since the integer m may be chosen freely.

Remark 1.13. Although it is not used elsewhere in the paper, in Section 11 we prove a result in the spirit of Theorem 1.11 for maps $u : X \rightarrow Y$, where Y is a general metric space equipped with a sequence of coverings.

Analogy with light mappings in the topological category. We would like to point out that Theorems 1.10, 1.11 are analogous to certain results for topological spaces.

Recall that a continuous map $f : X \rightarrow Y$ is light (respectively discrete, monotone) if the point inverses $\{f^{-1}(y)\}_{y \in Y}$ are totally disconnected (respectively discrete, connected). If X is a compact metrizable space, then X has topological dimension $\leq n$ if and only if there is a light map $X \rightarrow \mathbb{R}^n$; one implication comes from the fact that closed light maps do not decrease topological dimension [Eng95, Theorem 1.12.4], and the other follows from a Baire category argument.

One may consider versions of light mappings in the Lipschitz category. One possibility is the notion appearing in Theorems 1.7 and 1.11:

Definition 1.14. A Lipschitz map $f : X \rightarrow Y$ between metric spaces is **Lipschitz light** if there is a $C \in (0, \infty)$ such that for every bounded subset $W \subset Y$, the $\text{diam}(W)$ -components of $f^{-1}(W)$ have diameter $\leq C \cdot \text{diam}(W)$.

The analog with the topological case then leads to:

Definition 1.15. A metric space X has **Lipschitz dimension** $\leq n$ iff there is a Lipschitz light map from $X \rightarrow \mathbb{R}^n$ where \mathbb{R}^n has the usual metric.

With this definition, Theorems 1.7 and 1.11 become results about metric spaces of Lipschitz dimension ≤ 1 .

To carry the topological analogy further, we note that if $f : X \rightarrow Y$ is a light map between metric spaces and X is compact, then [Dyc74, DU97], in a variation on monotone-light factorization, showed that there is an inverse system

$$Y \longleftarrow X_1 \longleftarrow \dots \longleftarrow X_k \longleftarrow \dots$$

and a compatible family of mappings $\{g_k : X \rightarrow X_k\}$ such that:

- The projections $X_k \leftarrow X_{k+1}$ are discrete.
- g_k gives a factorization of f :

$$Y \longleftarrow X_1 \longleftarrow \dots \longleftarrow X_k \xleftarrow{g_k} X.$$

- The point inverses of g_k have diameter $\leq \Delta_k$, where $\Delta_k \rightarrow 0$ as $k \rightarrow \infty$.
- $\{g_k\}$ induces a homeomorphism $g_\infty : X \rightarrow X_\infty$, where X_∞ is the inverse limit X_∞ of the system $\{X_k\}$.

Making allowances for the difference between the Lipschitz and topological categories, this compares well with Theorem 1.11.

Embeddability and nonembeddability of inverse limits in Banach spaces. Theorem 1.11 reduces the proof of Theorem 1.7 (and also Theorem 1.1) to:

Theorem 1.16. *Let $\{X_i\}_{i \in \mathbb{Z}}$ be an admissible inverse system, and m be the parameter in Definition 1.8. There is a constant $L = L(m) \in (0, 1)$ and a 1-Lipschitz map $f : X_\infty \rightarrow L_1$ such that for all $x, y \in X_\infty$,*

$$\|f(x) - f(y)\|_{L_1} \geq L^{-1} \bar{d}_\infty(x, y).$$

In a forthcoming paper [CKa], we show that if one imposes additional conditions on an admissible inverse system $\{X_i\}$, the inverse limit X_∞ will carry a doubling measure μ which satisfies a Poincaré inequality, such that for μ a.e. $x \in X_\infty$, the tangent space $T_x X_\infty$ (in the sense of [Che99]) is 1-dimensional. The results apply to Examples 1.2 and 1.4. Moreover, in these two examples – and typically for the spaces studied in [CKa] – the Gromov-Hausdorff tangent cones at almost every point will not be bilipschitz homeomorphic to \mathbb{R} . The non-embedding result of [CK09] then implies that such spaces do not bilipschitz embed in Banach spaces which satisfy the Radon-Nikodym property. Combining this with Theorem 1.16, we therefore obtain a large class of examples of doubling spaces which embed in L_1 , but not in any Banach space satisfying the Radon-Nikodym property, cf. Corollary 1.5.

Monotone geodesics. Suppose $\{X_i\}_{i \in \mathbb{Z}}$ is an admissible inverse system, and $\phi : X_\infty \rightarrow \mathbb{R}$ is as in Theorem 1.10. Then ϕ picks out a distinguished class of paths, namely the paths $\gamma : I \rightarrow X_\infty$ such that the composition $\phi \circ \gamma : I \rightarrow \mathbb{R}$ is a homeomorphism onto its image, i.e. $\phi \circ \gamma : I \rightarrow \mathbb{R}$ is a monotone. (This is equivalent to saying that the projection $\pi_i \circ \gamma : I \rightarrow X_i$ is either direction preserving or direction reversing, with respect to the direction on X_i .) It is not difficult to see that such a path γ is a geodesic in X_∞ ; see Section 2. We call the image of such a path γ a **monotone geodesic segment** (respectively **monotone ray**, **monotone geodesic**) if the image $\phi \circ \gamma(I) \subset \mathbb{R}$ is a segment (respectively is a ray, is all of \mathbb{R}). Monotone geodesics and related structures play an important role in the proof of Theorem 1.16. In fact, the proof of Theorem 1.16 produces an embedding $f : X_\infty \rightarrow L_1$ with the additional property that it maps monotone geodesic segments in X_∞ isometrically to geodesic segments in L_1 .

Now suppose $u : X \rightarrow \mathbb{R}$ is as in Theorem 1.11. As above, one obtains a distinguished family of paths $\gamma : I \rightarrow X$, those for which $u \circ \gamma : I \rightarrow \mathbb{R}$ is a homeomorphism onto its image. From the assumptions on u , it is

easy to see that u induces a bilipschitz homeomorphism from the image $\gamma(I) \subset X$ to the image $(u \circ \gamma)(I) \subset \mathbb{R}$, so $\gamma(I)$ is a bilipschitz embedded path. We call the images of such paths **monotone**, although they need not be geodesics. If $f_\infty : X \rightarrow X_\infty$ is a homeomorphism provided by Theorem 1.11, then f_∞ maps monotone paths in X to monotone segments/rays/geodesics in X_∞ because $\phi \circ f_\infty = u$. Therefore, by combining Theorems 1.11 and 1.16, it follows that the embedding in Theorem 1.7 can be chosen to map monotone paths in X to geodesics in L_1 .


Discussion of the proof of Theorem 1.16. Before entering into the construction, we recall some observations from [CKb, CK10a, CK10b] which motivate the setup, and also indicate the delicacy of the embedding problem.

Let $\{X_i\}_{i \in \mathbb{Z}}$ be an admissible inverse system.

Suppose $f : X_\infty \rightarrow L_1$ is an L -bilipschitz embedding, and that X_∞ satisfies a Poincaré inequality with respect to a doubling measure μ (e.g. Examples 1.2 and 1.4). Then there is a version of Kirchheim's metric differentiation theorem [Kir94], which implies that for almost every $p \in X_\infty$, if one rescales the map f and passes to a limit, one obtains an L -bilipschitz embedding $f_\infty : Z \rightarrow L_1$, where Z is a Gromov-Hausdorff tangent space of X_∞ , such that $(f_\infty)|_\gamma : \gamma \rightarrow L_1$ is a constant speed geodesic for every $\gamma \subset Z$ which arises as a limit of (a sequence of rescaled) monotone geodesics in X_∞ . When X_∞ is self-similar, as in Examples 1.2 and 1.4, then Z contains copies of X_∞ , and one concludes that X_∞ itself has an L -bilipschitz embedding $X_\infty \rightarrow L_1$ which restricts to a constant speed geodesic embedding on each monotone geodesic $\gamma \subset X_\infty$. In view of this, and the fact that any bilipschitz embedding is constrained to have this behavior infinitesimally, our construction has been chosen so as to automatically satisfy the constraint, i.e. it generates maps which restrict to isometric embeddings on monotone geodesics.

By [Ass80, DL97, CK10a], producing a bilipschitz embedding $f : X_\infty \rightarrow L_1$ is equivalent to showing that distance function \bar{d}_∞ is comparable to a cut metric d_Σ , i.e. a distance function d_Σ on X_∞ which is a superposition of elementary cut metrics. Informally speaking this means that

$$d_\Sigma = \int_{2^{X_\infty}} d_E d\Sigma(E) .$$



Some children of E




FIGURE 2.

where Σ is a cut measure on the subsets of X_∞ , and d_E is the elementary cut (pseudo)metric associated with a subset $E \subset X_\infty$:

$$d_E(x_1, x_2) = |\chi_E(x_1) - \chi_E(x_2)|.$$

If $f : X_\infty \rightarrow L_1$ restricts to an isometric embedding $f|_\gamma : \gamma \rightarrow L_1$ for every monotone geodesic $\gamma \subset X_\infty$, then one finds (informally speaking) that the cut measure Σ is supported on subsets $E \subset X_\infty$ with the property that for every monotone geodesic $\gamma \subset X_\infty$, the characteristic function χ_E restricts to a monotone function on γ , or equivalently, that

the the intersections $E \cap \gamma$ and $(X_\infty \setminus E) \cap \gamma$ are both connected. We call such subsets **monotone**.

For simplicity we restrict the rest of our discussion to the case when $X_0 \simeq \mathbb{R}$. The reader may find it helpful to keep Example 1.2 in mind (modified with $X_i \simeq \mathbb{R}$ for $i < 0$ as indicated earlier).

Motivated by the above observations, the approach taken in the paper is to obtain the cut metric d_Σ as a limit of a sequence of cut metrics $\{d_{\Sigma'_i}\}_{i \geq 0}$, where Σ'_i is a cut measure on X_i supported on monotone subsets. For technical reasons, we choose Σ'_i so that every monotone subset E in the support of Σ'_i is a subcomplex of X'_i (see Definition 1.8), and E is precisely the set of points $x \in X_i$ such that there is a monotone geodesic $c : [0, 1] \rightarrow X_i$ where $\pi_0^i \circ c$ is increasing, $c(0) = x$, and $c(1)$ lies in the boundary of E ; thus one may think of E as the set of points “lying to the left” of the boundary ∂E .

We construct the sequence $\{\Sigma'_i\}$ inductively as follows. The cut measure Σ'_0 is the atomic measure which assigns mass $\frac{1}{m}$ to each monotone subset of the form $(-\infty, v]$, where v is vertex of $X'_0 \simeq \mathbb{R}$. Inductively we construct Σ'_{i+1} from Σ'_i by a diffusion process. For every monotone set $E \subset X_i$ in the support of Σ'_i , we take the Σ'_i -measure living on E , and redistribute it over a family of monotone sets $E' \subset X_{i+1}$, called the **children** of E . The children of $E \subset X_i$ are monotone sets $E' \subset X_{i+1}$ obtained from the inverse image $\pi_i^{-1}(E)$ by modifying the boundary locally: for each vertex v of X_{i+1} lying in the boundary of $\pi_i^{-1}(E)$, we move the boundary within the open star of v . An example of this local modification procedure is depicted in Figure 2, where $m = 4$.

The remainder of the proof involves a series of estimates on the cut measures Σ'_i and cut metrics $d_{\Sigma'_i}$, which are proved by induction on i using the form of the diffusion process, see Section 7. One shows that the sequence of pseudo-metrics $\{\rho_i = d_{\Sigma'_i} \circ \pi_i^\infty\}$ on X_∞ converges geometrically to a distance function which will be the cut metric d_Σ for a cut measure Σ on X_∞ . To prove that d_Σ is comparable to d_∞ , the idea is to show (by induction) that the cut metric $d_{\Sigma'_i}$ resolves pairs of points $x_1, x_2 \in X_i$ whose separation is $> C m^{-i}$.

Organization of the paper. In Section 2 we collect notation and establish some basic properties of admissible inverse systems. Theorem 1.10 is proved in Section 2.3. Section 3 considers a special class of admissible inverse systems which come with natural metrics, e.g. Examples 1.2, 1.4. In Section 4 we prove Theorem 1.11. Sections 5–9

give the proof of Theorem 1.16. A special case of Theorem 1.16 is introduced in Section 5. In Section 6 we begin the proof of the special case by developing the structure of slices and associated slice measures, which are closely related to the monotone sets in the above discussion of the proof of Theorem 1.16. Section 7 obtains estimates on the slice measures which are needed for the embedding theorem. Section 8 completes the proof of Theorem 1.16 in the special case introduced in Section 5. Section 9 completes the proof in the general case. Section 10 shows that the space in Example 1.4 is bilipschitz homeomorphic to a Laakso space from [Laa00]. In Section 11 we consider a generalization of Theorem 1.11 to maps $u : X \rightarrow Y$, where Y is a general metric space equipped with a sequence of coverings.

We refer the reader to the beginnings of the individual sections for more detailed descriptions of their contents.

2. NOTATION AND PRELIMINARIES

In this section $\{X_i\}_{i \in \mathbb{Z}}$ will be an admissible inverse system, and m will be the parameter appearing in Definition 1.8.

2.1. Subdivisions, stars, and trimmed stars. Let Z be a graph. Let $Z^{(k)}$ denote the k -fold iterated subdivision of Z , where each iteration subdivides every edge into m subedges, and let $Z' = Z^{(1)}$.

If v is a vertex of a graph \mathcal{G} , then $\text{St}(v, \mathcal{G})$ and $\text{St}^o(v, \mathcal{G})$ denote the closed and open stars of v , respectively.

Definition 2.1. Let Z be a graph, and $v \in Z$ be a vertex. The **trimmed star of v in Z** is the union of the edges of Z' which lie in the open star $\text{St}^o(v, Z)$, or alternately, the union of the edge paths in Z' starting at v , with $(m - 1)$ edges. We denote the trimmed star by $\text{TSt}(v, Z)$. We will only use this when $Z = X_i$ or $Z = X'_i$ below.

Note that if v is a vertex of X_i , then $\text{TSt}(v, X_i)$ is also the closed ball $\overline{B(v, \frac{m-1}{m} \cdot m^{-i})} \subset X_i$ with respect to the path metric d_i .

2.2. Basic properties of admissible inverse systems and the distance function \bar{d}_∞ . Let $\{X_i\}_{i \in \mathbb{Z}}$ be an admissible inverse system with inverse limit X_∞ . For every i , we let V_i be the vertex set of X_i ,

and V'_i be the vertex set of X'_i . For all $j \geq i$, let $\pi_i^j : X_j \rightarrow X_i$ be the composition $\pi_{j-1} \circ \dots \circ \pi_i$. Then $\pi_i^j : X_j \rightarrow X_i^{(j-i)}$ is simplicial and restricts to an isomorphism on each edge. It is also 1-Lipschitz with respect to the respective path metrics d_j and d_i .

Lemma 2.2. *For every $x, x' \in X_\infty$ there exist $i \in \mathbb{Z}$, $v \in V_i$ such that $x, x' \in (\pi_i^\infty)^{-1}(\text{St}(v, X_i))$.*

Proof. By (3) of Definition 1.8, there is a $j \in \mathbb{Z}$ such that $\pi_j(x), \pi_j(x')$ are contained in the same connected component of X_j . If $\gamma \subset X_j$ is a path from $\pi_j(x)$ to $\pi_j(x')$ with d_j -length N , then for all $i \leq j$, the projection $\pi_i^j(\gamma)$ is a path in $X_i^{(j-i)}$ with d_i -length $\leq N$. Therefore if $N < m^{-i}$ then $\pi_i^j(\gamma)$ will be contained in $\text{St}(v, X_i)$ for some $v \in V_i$. \square

Suppose \hat{d} is a pseudo-distance on X_∞ with the property that

$$\text{diam}_{\hat{d}}((\pi_j^\infty)^{-1}(\text{St}(v, X_j))) \leq 2m^{-j}$$

for all $j \in \mathbb{Z}$, $v \in V_j$. Then for every $x, x' \in X_\infty$, we have $\hat{d}(x, x') \leq 2m^{-i}$, where $i \in \mathbb{Z}$ is as in Lemma 2.2. It follows that the supremum \bar{d}_∞ of all such pseudo-distance functions takes finite values, i.e. is a well-defined pseudo-distance function.

Lemma 2.3 (Alternate definition of \bar{d}_∞). *Suppose $x, x' \in X_\infty$. Then $\bar{d}_\infty(x, x')$ is the infimum of the sums $\sum_{k=1}^n 2m^{-i_k}$, such that there exists a finite sequence*

$$x = x_0, \dots, x_n = x' \in X_\infty$$

where $\{\pi_{i_k}^\infty(x_{k-1}), \pi_{i_k}^\infty(x_k)\}$ is contained in the closed star $\text{St}(v, X_{i_k})$ for some vertex v of X_{i_k} , for every $k \in \{1, \dots, n\}$.

Proof. Let $\hat{d}_\infty(x, x') \in [0, \infty]$ be the infimum defined above. By Lemma 2.2 the infimum will be taken over a nonempty set of sequences, and so $\hat{d}_\infty(x, x') \in [0, \infty)$. It follows that \hat{d}_∞ is a well-defined pseudo-distance satisfying the condition that $\text{diam}_{\hat{d}_\infty}((\pi_i^\infty)^{-1}(\text{St}(v, X_i))) \leq 2m^{-i}$ for every $v \in V_i$, $i \in \mathbb{Z}$. Therefore $\hat{d}_\infty \leq \bar{d}_\infty$ from the definition of \bar{d}_∞ . On the other hand the definition of \bar{d}_∞ and the triangle inequality imply $\bar{d}_\infty \leq \hat{d}_\infty$. \square

Lemma 2.4. *Suppose $x, x' \in X_\infty$.*

- (1) *If $\bar{d}_\infty(x, x') \leq m^{-j}$ for some $j \in \mathbb{Z}$, then $\pi_j(x), \pi_j(x')$ belong to $\text{St}(v, X_j)$ for some $v \in V_j$.*

- (2) If $x \in X_\infty$ and $r \leq \frac{(m-2)}{m}m^{-j}$ for some $j \geq 0$, then $\pi_j(B(x, r))$ is contained in the trimmed star $\text{TSt}(v, X_j)$ for some $v \in V_j$.

Proof. (1). Pick $\epsilon > 0$. Since $\bar{d}_\infty(x, x') \leq m^{-j}$, there is a sequence $x = y_0, \dots, y_k = x' \in X_\infty$, where for $\ell \in \{1, \dots, k\}$, the points $y_{\ell-1}, y_\ell$ lie in $\pi_{i_\ell}^{-1}(\text{St}(v_\ell, X_{i_\ell}))$, $v_\ell \in V_{i_\ell}$, and

$$\sum_{\ell} 2m^{-i_\ell} \leq m^{-j} + \epsilon.$$

Taking $\epsilon < m^{-j}$, we may assume that $i_\ell \geq j$ for all $\ell \in \{1, \dots, k\}$. Since $\pi_{i_\ell}(y_{\ell-1}), \pi_{i_\ell}(y_\ell) \in \text{St}(v_\ell, X_{i_\ell})$, there is a path from $\pi_{i_\ell}(y_{\ell-1})$ to $\pi_{i_\ell}(y_\ell)$ in X_{i_ℓ} of d_{i_ℓ} -length $\leq 2m^{-i_\ell}$. Since $\pi_j^{i_\ell} : (X_{i_\ell}, d_{i_\ell}) \rightarrow (X_j, d_j)$ is 1-Lipschitz for all j , we get that there is a path from $\pi_j(x)$ to $\pi_j(x')$ in X_j with d_j -length at most

$$\sum_{\ell} 2m^{-i_\ell} < m^{-j} + \epsilon.$$

As ϵ is arbitrary, $\pi_j(x)$ and $\pi_j(x')$ lie in $\text{St}(v, X_j)$ for some $j \in V_j$.

(2). The proof is similar to (1). \square

Corollary 2.5. \bar{d}_∞ is a distance function on X_∞ .

Proof. Suppose $x, x' \in X_\infty$, $\bar{d}_\infty(x, x') = 0$, and $i \in \mathbb{Z}$. By Lemma 2.4, for all $j \geq i$ the set $\{\pi_j^\infty(x), \pi_j^\infty(x')\}$ is contained in the star of some vertex $v \in V_j$. Since $\pi_i^j : (X_j, d_j) \rightarrow (X_i, d_i)$ is 1-Lipschitz, it follows that $\{\pi_i^\infty(x), \pi_i^\infty(x')\}$ is contained in a set of d_i -diameter $\leq 2m^{-j}$. Since j is arbitrary, this means that $\pi_i^\infty(x) = \pi_i^\infty(x')$. \square

The following is a sharper statement:

Lemma 2.6. Suppose $x_1, x_2 \in X_\infty$ are distinct points. Let j be the minimum of the indices $k \in \mathbb{Z}$ such that $\{\pi_k(x_1), \pi_k(x_2)\}$ is not contained in the trimmed star $\text{TSt}(v, X_k)$ for any $v \in V_k$. Then

$$(2.7) \quad \frac{(m-2)}{m}m^{-j} < \bar{d}_\infty(x_1, x_2) \leq 2m^{-(j-1)}.$$

Proof. The first inequality follows immediately from Lemma 2.4. By the choice of j , there is a vertex $v \in V_{j-1}$ such that

$$\{\pi_{j-1}(x_1), \pi_{j-1}(x_2)\} \subset \text{TSt}(v, X_{j-1}) \subset \text{St}(v, X_{j-1}),$$

so $\bar{d}_\infty(x_1, x_2) \leq 2m^{-(j-1)}$ by Definition 1.9. \square

2.3. A canonical map from the inverse limit to \mathbb{R} . The next theorem contains Theorem 1.10.

Theorem 2.8. *Suppose $\{X_i\}$ is an admissible inverse system.*

- (1) *There is a compatible system of direction preserving maps $\phi_i : X_i \rightarrow \mathbb{R}$, such that for every i , the restriction of ϕ_i to any edge $e \subset X_i$ is a linear map onto a segment of length m^{-i} . In particular, ϕ_i is 1-Lipschitz with respect to d_i .*
- (2) *The system of maps $\{\phi_i : X_i \rightarrow \mathbb{R}\}$ is unique up to post-composition with translation.*
- (3) *If $\phi : (X_\infty, \bar{d}_\infty) \rightarrow \mathbb{R}$ is the map induced by $\{\phi_i\}$, then ϕ is 1-Lipschitz, and for every interval $I \subset \mathbb{R}$, the $\text{diam}(I)$ -components of $\phi^{-1}(I)$ have diameter at most $8m \cdot \text{diam}(I)$.*

Proof. (1). Let X_∞ denote the direct limit of the system $\{X_i\}$, i.e. X_∞ is the disjoint union $\sqcup_{i \in \mathbb{Z}} X_i$ modulo the equivalence relation that $X_i \ni x \sim x' \in X_j$ if and only if there is a $k \leq \min(i, j)$ such that $\pi_k^i(x) = \pi_k^j(x')$. For every $i \in \mathbb{Z}$ there is a canonical projection map $\pi_\infty^i : X_i \rightarrow X_\infty$.

If $k \in \mathbb{Z}$, then for all $i \leq k$ let $X_i^{(k-i)}$ denote the $(k-i)$ -fold iterated subdivision of X_i , as in Section 2.1. Thus $\pi_i^j : X_j^{(k-j)} \rightarrow X_i^{(k-i)}$ is simplicial for all $i \leq j \leq k$, and restricts to a direction-preserving isomorphism on each edge of $X_j^{(k-j)}$. Therefore the direct limit X_∞ inherits a directed graph structure, which we denote $X_\infty^{(k-\infty)}$, and for all $i \leq k$, the projection map $\pi_\infty^i : X_i^{(k-i)} \rightarrow X_\infty^{(k-\infty)}$ is simplicial, and a directed isomorphism on each edge of $X_i^{(k-i)}$. Condition (3) of the definition of admissible systems implies that $X_\infty^{(k-\infty)}$ is connected.

Note also that for all $k \leq l$, the graph $X_\infty^{(l-\infty)}$ is canonically isomorphic to $(X_\infty^{(k-\infty)})^{(l-k)}$. In particular, if v, v' are distinct vertices of $X_\infty^{(k-\infty)}$, then their combinatorial distance in $X_\infty^{(l-\infty)}$ is at least m^{l-k} ; moreover every vertex of $X_\infty^{(l-\infty)}$ which is not a vertex of $X_\infty^{(k-\infty)}$ must have valence 2, since it corresponds to an interior point of an edge of $X_\infty^{(k-\infty)}$. It follows that $X_\infty^{(k-\infty)}$ can contain at most one vertex v which has valence $\neq 2$. Thus $X_\infty^{(k-\infty)}$ is either isomorphic to \mathbb{R} with the standard subdivision, or to the union of a single vertex v with a (possibly empty) collection of standard rays, each of which is direction-preserving isomorphic to either $(-\infty, 0]$ or $[0, \infty)$ with the standard subdivision. In either case, there is clearly a direction preserving simplicial map

$X_{-\infty}^{(k-\infty)} \rightarrow \mathbb{R}$ which is an isomorphism on each edge of $X_{-\infty}^{(k-\infty)}$. Pre-composing this with the projection maps $X_{\infty} \rightarrow X_k \rightarrow X_{-\infty}$ gives the desired maps ϕ_i .

(2). Any such system $\{\phi_i : X_i \rightarrow \mathbb{R}\}$ induces a map $\phi_{-\infty} : X_{-\infty} \rightarrow \mathbb{R}$, which for all $k \in \mathbb{Z}$ restricts to a direction preserving isomorphism on every edge of $X_{-\infty}^{(k-\infty)}$. From the description of $X_{-\infty}^{(k-\infty)}$, the map $\phi_{-\infty}$ is unique up to post-composition with a translation.

(3). If $x, x' \in X_{\infty}$ and $\{\pi_i(x), \pi_i(x')\} \subset \text{St}(v, X_i)$ for some $i \in \mathbb{Z}$, $v \in V_i$, then by (1) $\{\phi(x), \phi(x')\}$ is contained in the union of two intervals of length m^{-i} in \mathbb{R} , and therefore $d(\phi(x), \phi(x')) \leq 2m^{-i}$. By the definition of \bar{d}_{∞} , this implies that $d(\phi(x), \phi(x')) \leq \bar{d}_{\infty}(x, x')$ for all $x, x' \in X_{\infty}$, i.e. ϕ is 1-Lipschitz.

From the construction of the map $X_{-\infty} \rightarrow \mathbb{R}$, there exists a sequence $\{Y_i\}_{i \in \mathbb{Z}}$ of subdivisions of \mathbb{R} , such that $Y_{i+1} = Y_i^{(1)}$, and $\phi_i : X_i \rightarrow \mathbb{R} \simeq Y_i$ is simplicial and restricts to an isomorphism on every edge of X_i .

Now suppose $I \subset \mathbb{R}$ is an interval, and choose $i \in \mathbb{Z}$ such that $\text{diam}(I) \in [\frac{m^{-(i+1)}}{4}, \frac{m^{-i}}{4})$. Then there is a vertex $v \in Y_i$ such that $I \subset \text{St}(v, Y_i)$ and $\text{dist}(I, \mathbb{R} \setminus \text{St}(v, Y_i)) > \text{diam}(I)$. Pick $x, x' \in \phi^{-1}(I)$ which lie in the same $\text{diam}(I)$ -component of $\phi^{-1}(I) \subset X_{\infty}$, so there is a $\text{diam}(I)$ -path $x = x_0, \dots, x_k = x'$ in X_{∞} . For each $\ell \in \{1, \dots, k\}$ and every $\epsilon > 0$, there is a path γ_{ℓ} in X_i which joins $\pi_i(x_{\ell-1})$ to $\pi_i(x_{\ell})$ such that $\text{length}(\phi_i \circ \gamma_{\ell}) \leq \text{diam}(I) + \epsilon$. When ϵ is sufficiently small we get $\gamma_{\ell} \subset \phi_i^{-1}(\text{St}^o(v, Y_i))$ because $\phi_i \circ \gamma_{\ell}$ has endpoints in I . Therefore $\pi_i(x), \pi_i(x')$ lie in the same path component of $\phi_i^{-1}(\text{St}^o(v, Y_i))$, which implies that they lie in $\text{St}(v, X_i)$ for some vertex $v \in V_i$. Hence $\bar{d}_{\infty}(x, x') \leq 2m^{-i} \leq 8m \cdot \text{diam}(I)$.

□

2.4. Directed paths, a partial ordering, and monotone paths.

Suppose $\{X_i\}$ is an admissible system, and $\{\phi_i : X_i \rightarrow \mathbb{R}\}$ is a system of maps as in Theorem 2.8.

Definition 2.9. A **directed path** in X_i is a path $\gamma : I \rightarrow X_i$ which is locally injective, and direction preserving (w.r.t. the usual direction on I). A **directed path** in X_{∞} is a path $\gamma : I \rightarrow X_{\infty}$ such that $\pi_i \circ \gamma : I \rightarrow X_i$ is directed for all $i \in \mathbb{Z}$.

If $\gamma : I \rightarrow X_j$ is a directed path in X_j , then $\phi_j \circ \gamma$ is a directed path in \mathbb{R} , and hence it is embedded, and has the same length as γ . Therefore X_∞ and the X_j 's do not contain directed loops. Furthermore, it follows that $\pi_i^j \circ \gamma$ is a d_i -geodesic in X_i for all $i \leq j$.

Definition 2.10 (Partial order). We define a binary relation on X_i , for $i \in \mathbb{Z} \cup \{\infty\}$ by declaring that $x \preceq y$ if there is a (possibly trivial) directed path from x to y . This defines a partial order on X_i since X_i contains no directed loops. As usual, $x \prec y$ means that $x \preceq y$ and $x \neq y$.

Since the projections $\pi_i^j : X_j \rightarrow X_i$ are direction-preserving, they are order preserving for all $i \leq j$, as is the projection map $\pi_i^\infty : X_\infty \rightarrow X_i$.

Lemma 2.11. *Suppose $\gamma : I \rightarrow X_\infty$ is a continuous map. The following are equivalent:*

- (1) γ is a directed geodesic, i.e. $\text{length}(\gamma) = d(\gamma(0), \gamma(1))$.
- (2) γ is a directed path.
- (3) $\pi_i \circ \gamma$ is a directed path for all i .
- (4) $\phi \circ \gamma : [0, 1] \rightarrow \mathbb{R}$ is a directed path.

Proof. (1) \implies (2) \implies (3) \implies (4) is clear.

(4) \implies (3) follows from the fact that $\phi_i : X_i \rightarrow \mathbb{R}$ restricts to a direction preserving isomorphism on every edge of X_i .

(3) \implies (1). For all $i \in \mathbb{Z}$, let $\gamma_i \subset X_i$ be the union of the edges whose interiors intersect the image of $\pi_i \circ \gamma$. Then γ_i is a directed edge path in X_i , and hence $\phi_i \circ \gamma_i$ is a directed edge path in \mathbb{R} with the same number of edges. Therefore γ_i has $< 2 + m^i(d(\phi(\gamma(0)), \phi(\gamma(1))))$ edges. Since the vertices of γ_i belong to the image of $\pi_i : X_\infty \rightarrow X_i$, by the definition of \bar{d}_∞ , we have $\bar{d}_\infty(\gamma(0), \gamma(1)) \leq 3m^{-i} + d(\phi(\gamma(0)), \phi(\gamma(1)))$. Since i is arbitrary we get $\bar{d}_\infty(\gamma(0), \gamma(1)) \leq d(\phi(\gamma(0)), \phi(\gamma(1)))$, and Theorem 2.8(3) gives equality. This holds for all subpaths of γ as well, so γ is a geodesic. \square

Definition 2.12. A **monotone geodesic segment in X_i** is the image of a directed isometric embedding $\gamma : [a, b] \rightarrow X_i$; a **monotone geodesic in X_i** is the image of a directed isometric embedding $\mathbb{R} \rightarrow (X_i, d_i)$. A **monotone geodesic segment in X_∞** is (the image of) a path $\gamma : I \rightarrow X_\infty$ satisfying any of the conditions of the lemma.

A **monotone geodesic** is (the image of) a directed isometric embedding $\mathbb{R} \rightarrow X_\infty$, or equivalently, a geodesic $\gamma \subset X_\infty$ which projects isometrically under $\phi : X_\infty \rightarrow \mathbb{R}$ onto \mathbb{R} .

Monotone geodesics lead to monotone sets:

Definition 2.13. A subset $E \subset X_i$, $i \in \mathbb{Z} \cup \{\infty\}$, is **monotone** if the characteristic function χ_E restricts to a monotone function on any monotone geodesic $\gamma \subset X_i$ (i.e. $\gamma \cap E$ and $\gamma \setminus E$ are both connected subsets of γ).

3. INVERSE SYSTEMS OF GRAPHS WITH PATH METRICS

For some admissible inverse systems, such as Examples 1.2, 1.4, the path metrics d_i induce a length structure on the inverse limit which is comparable to \bar{d}_∞ . We discuss this special class here, comparing the length metric with the metric \bar{d}_∞ defined earlier.

In this section we assume that $\{X_i\}$ is an admissible inverse system satisfying two additional conditions:

- (a) $\pi_i : X_{i+1} \rightarrow X'_i$ is an open map for all $i \in \mathbb{Z}$.
- (b) There is a $\theta \in \mathbb{N}$ such that for every $i \in \mathbb{Z}$, $v \in V_i$, $w, w' \in \pi_i^{-1}(v)$, there is an edge path in X_{i+1} with at most θ edges, which joins w and w' .

Both conditions obviously hold in Examples 1.2, 1.4.

Lemma 3.1 (Path lifting). *Suppose $0 \leq i < j$, $c : [a, b] \rightarrow X_i$ is a path, and $v \in (\pi_i^j)^{-1}(c(a))$. Then there is a path $\hat{c} : [a, b] \rightarrow X_j$ such that*

- \hat{c} is a lift of c : $\pi_i^j \circ \hat{c} = c$.
- $\hat{c}(a) = v$.

Proof. If c is piecewise linear, and linear on each subinterval of the partition $a = t_0 < \dots < t_k = b$, then the existence of \hat{c} follows by induction on k , because of condition (a) above. The general case follows by approximation. \square

Lemma 3.2.

- (1) X_i is connected for all i .
- (2) $\pi_i^j : X_j \rightarrow X_i$ and $\pi_i^\infty : X_\infty \rightarrow X_i$ are surjective for all $i \leq j$.

Proof. (1). By Lemma 3.1 and condition (b), if $x_1, x_2 \in X_i$ lie in the same connected component of X_i , then any $x'_1 \in \pi_i^{-1}(x_1)$, $x'_2 \in \pi_i^{-1}(x_2)$ lie in the same connected component of X_{i+1} . Iterating this, we get that $(\pi_i^j)^{-1}(x_1) \cup (\pi_i^j)^{-1}(x_2) \subset X_j$ is contained in a single component of X_j . Now for every $j \in \mathbb{Z}$, $\hat{x}_1, \hat{x}_2 \in X_j$, by Definition 1.8 (3) there is an $i \leq j$ such that $x_1 = \pi_i(\hat{x}_1)$, $x_2 = \pi_i(\hat{x}_2)$ lie in the same connected component of X_i ; therefore \hat{x}_1, \hat{x}_2 lie in the same component of X_j .

(2). $\pi_i : X_{i+1} \rightarrow X_i$ is open by condition (a), X_i is connected by (1), and X_{i+1} is nonempty by Definition 1.8 (1). Therefore $\pi_i : X_{i+1} \rightarrow X_i$ is surjective. It follows that $\pi_i^\infty : X_\infty \rightarrow X_i$ is surjective as well. \square

Note that $\pi_i : (X_{i+1}, d_{i+1}) \rightarrow (X_i, d_i)$ is a 1-Lipschitz map by Definition 1.8(2).

Lemma 3.3.

(1) For all $i \in \mathbb{Z}$, and every $x_1, x_2 \in X_i$, $x'_1, x'_2 \in X_{i+1}$ with $\pi_i(x'_k) = x_k$, we have

$$d_{i+1}(x'_1, x'_2) \leq d_i(x_1, x_2) + 2m^{-i} + \theta \cdot m^{-(i+1)}.$$

(2) If $i < j$, then for every $x_1, x_2 \in X_i$, $x'_1, x'_2 \in X_j$ with $\pi_i(x'_k) = x_k$, we have

$$(3.4) \quad d_j(x'_1, x'_2) \leq d_i(x_1, x_2) + (2m + \theta) \cdot \left(\frac{m^{-i} - m^{-j}}{m - 1} \right).$$

Proof. (1). Let $\gamma : I \rightarrow X_i$ be a path of length at most $d_i(x_1, x_2) + m^{-i}$ which joins x_1 to x_2 , and then continues to some vertex $v_2 \in V_i$. By Lemma 3.1 there is a lift $\gamma' : I \rightarrow X_{i+1}$ starting at x'_1 , and clearly $\text{length}(\gamma') = \text{length}(\gamma)$. Since x'_2 has distance $< m^{-i}$ from $\pi_i^{-1}(v_2)$, (1) follows from condition (b).

(2). This follows by iterating (1). \square

As a consequence of Lemma 3.3, the sequence of (pseudo)distance functions $\{d_i \circ \pi_i^\infty : X_\infty \times X_\infty \rightarrow [0, \infty)\}_{i \geq 0}$ converges geometrically to a distance function d_∞ on X_∞ . Since π_i^∞ is surjective for all $i \geq 0$, the lemma also implies that $\{\pi_i^\infty : (X_\infty, d_\infty) \rightarrow (X_i, d_i)\}_{i \geq 0}$ is a sequence of Gromov-Hausdorff approximations, so (X_i, d_i) converges to (X_∞, d_∞) in the Gromov-Hausdorff topology.

Lemma 3.5.

- (1) $\bar{d}_\infty \leq d_\infty$, with equality on monotone geodesic segments.
 (2) $d_\infty \leq \frac{2m+\theta}{2(m-1)} \cdot \bar{d}_\infty$.

Proof. (1). Suppose $x, x' \in X_\infty$ and for some $i \in \mathbb{Z}$ $\gamma : I \rightarrow X_i$ is a geodesic from $\pi_i(x)$ to $\pi_i(x')$. Then the image of γ is contained in a chain of at most $2 + \frac{d_\infty(x, x')}{2m^{-i}}$ stars in X_i . Since π_i^∞ is surjective, Definition 1.9 implies

$$\bar{d}_\infty(x, x') \leq d_\infty(x, x') + 2m^{-i}.$$

Thus $\bar{d}_\infty \leq d_\infty$.

If $x, x' \in X_\infty$ are joined by a directed path $\gamma : I \rightarrow X_\infty$, then

$$\begin{aligned} d_\infty(x, x') &\leq \text{length}(\gamma) = \text{length}(\pi_i \circ \gamma) \\ &= d(\phi(\gamma(0)), \phi(\gamma(1))) \leq \bar{d}_\infty(x, x'), \end{aligned}$$

where the last equality follows from Theorem 2.8(3).

(2). If $x, x' \in X_\infty$ and $\{\pi_i(x), \pi_i(x')\} \subset \text{St}(v, X_i)$ for some $i \in \mathbb{Z}$, $v \in V_i$, then $d_i(\pi_i(x), \pi_i(x')) \leq 2m^{-i}$ and so

$$d_\infty(x, x') \leq \frac{(2m + \theta)m^{-i}}{m - 1}$$

by Lemma 3.3. It follows from Lemma 2.3 that

$$d_\infty \leq \frac{(2m + \theta)}{2(m - 1)} \cdot \bar{d}_\infty.$$

□

Corollary 3.6. *If $\phi : X_\infty \rightarrow \mathbb{R}$ is the map from Theorem 2.8, then $\phi : (X_\infty, d_\infty) \rightarrow \mathbb{R}$ satisfies the hypotheses of Theorem 1.1.*

Proof. This follows from the previous Lemma and Theorem 2.8(3). □

4. REALIZING METRIC SPACES AS LIMITS OF ADMISSIBLE INVERSE SYSTEMS

In this section, we characterize metric spaces which are bilipschitz homeomorphic to inverse limits of admissible inverse systems, proving Theorem 1.11.

Suppose a metric space Z is bilipschitz equivalent to the inverse limit of an admissible inverse system. Evidently, if X_∞ is such an inverse limit, $\phi : X_\infty \rightarrow \mathbb{R}$ is as in Theorem 2.8, and $F : Z \rightarrow X_\infty$ is a bilipschitz homeomorphism, then the composition $u = \phi \circ F : Z \rightarrow \mathbb{R}$ has the property that for every interval $I \subset \mathbb{R}$, the $\text{diam}(I)$ -components of $u^{-1}(I)$ have diameter at most comparable to $\text{diam}(I)$, (see Theorem 2.8). In other words a necessary condition for a space to be bilipschitz homeomorphic to an inverse limit is the existence of a map satisfying the hypotheses of Theorem 1.11. Theorem 1.11 says that the existence of such a map is sufficient.

We now prove Theorem 1.11.

Fix $m \in \mathbb{N}$, $m \geq 2$, and let $u : X \rightarrow \mathbb{R}$ be as in the statement of the theorem.

Let $\{Y_i\}_{i \in \mathbb{Z}}$ be a sequence of subdivisions of \mathbb{R} , where:

- Y_i is a subdivision of \mathbb{R} into intervals of length m^{-i} for all $i \in \mathbb{Z}$.
- Y_{i+1} is a subdivision of Y_i for all $i \in \mathbb{Z}$.

We define a simplicial graph X_i as follows. The vertex set of X_i is the collection of pairs (v, U) where v is a vertex of Y_i and U is a m^{-i} -component of $u^{-1}(\text{St}(v, Y_i))$. Two distinct vertices $(v_1, U_1), (v_2, U_2) \in X_i$ span an edge iff $U_1 \cap U_2 \neq \emptyset$; note that this can only happen if v_1, v_2 are distinct adjacent vertices of Y_i .

We have an projection map $\phi_i : X_i \rightarrow Y_i$ which sends each vertex (v, U) of X_i to $v \in Y_i$, and is a linear isomorphism on each edge of X_i . If (\hat{v}, \hat{U}) is a vertex of X_{i+1} , there there will be a vertex (v, U) of X_i such that $\hat{U} \subset U$ and $\text{St}(\hat{v}, Y_{i+1}) \subset \text{St}(v, Y_i)$; there are at most two such vertices, and they will span an edge in X_i . Therefore we obtain a well-defined projection map $\pi_i : X_{i+1} \rightarrow X_i$ such that $\phi_i \circ \pi_i = \phi_{i+1}$, and which induces a simplicial map $\pi_i : X_{i+1} \rightarrow X'_i$.

We define $f_i : X \rightarrow X_i$ as follows. Suppose $z \in X$ and $u(z) \in \mathbb{R} \simeq Y_i$ belongs to the edge $e = \overline{v_1 v_2} = \text{St}(v_1, Y_i) \cap \text{St}(v_2, Y_i)$. Then z belongs to an m^{-i} -component of $\text{St}(v_j, Y_i)$ for $j \in \{1, 2\}$, and therefore these two components span an edge \hat{e} of X_i which is mapped isomorphically to e by ϕ_i . We define $f_i(z)$ to be $\phi_i^{-1}(u(z)) \cap \hat{e}$. The sequence $\{f_i\}_{i \in \mathbb{Z}}$ is clearly compatible, so we have a well-defined map $f_\infty : X \rightarrow X_\infty$.

Now suppose $z_1, z_2 \in X$ and for some $i \in \mathbb{Z}$ we have $d(z_1, z_2) \leq m^{-i}$. Then $\{u(z_1), u(z_2)\} \subset \text{St}(v, Y_i)$ for some vertex $v \in Y_i$, and z_1, z_2 lie in the same m^{-i} component of $u^{-1}(\text{St}(v, Y_i))$. Therefore $\{f_i(z_1), f_i(z_2)\}$

is contained in $\text{St}(\hat{v}, X_i)$ for some vertex \hat{v} of X_i . It follows that $\bar{d}_\infty(f_\infty(z_1), f_\infty(z_2)) \leq 2m^{-i}$, from the definition of \bar{d}_∞ . Thus f_∞ is Lipschitz.

On the other hand, if $z_1, z_2 \in X$ and $\bar{d}_\infty(f_\infty(z_1), f_\infty(z_2)) \leq m^{-i}$, then by Lemma 2.4, we have $\{f_j(z_1), f_j(z_2)\} \subset \text{St}(\hat{v}, X_j)$ for some vertex $\hat{v} \in X_j$, where $|i - j|$ is controlled. By the construction of f_j , this means that $\{z_1, z_2\}$ lie in an m^{-j} -component of $u^{-1}(\text{St}(v, Y_j))$ for $v = \phi_j(\hat{v}) \in Y_j$. By our assumption on u , this gives $d(z_1, z_2) \lesssim m^{-i}$. Thus f_∞ is L' -bilipschitz, where L' depends only on C and m .

5. A SPECIAL CASE OF THEOREM 1.16

Let $\{X_j\}$ be an admissible inverse system as in Definition 1.8.

Assumption 5.1. *We will temporarily assume that:*

- (1) π_i is finite-to-one for all $i \in \mathbb{Z}$.
- (2) $X_i \simeq \mathbb{R}$ and $\pi_i : X_{i+1} \rightarrow X_i'$ is an isomorphism for all $i \leq 0$.
- (3) For every $i \in \mathbb{Z}$, every vertex $v \in V_i$ has neighbors $v_\pm \in V_i$ with $v_- \prec v \prec v_+$. Equivalently, X_i is a union of (complete) monotone geodesics (see Definition 2.12).

In particular, (1) and (2) imply that X_i has finite valence for all i .

This extra assumption will be removed in Section 9, in order to complete the proof in the general case. We remark that it is possible to adapt all the material to the general setting, but this would impose a technical burden that is largely avoidable. Furthermore, Assumption 5.1 effectively covers many cases of interest, such as Examples 1.2 and 1.4.

6. SLICES AND THE ASSOCIATED MEASURES

Rather than working directly with monotone subsets as described in the introduction, we instead work with subsets which we call slices, which are sets of vertices which arise naturally as the boundaries of monotone subsets. A slice in $S \subset X_i$ gives rise to a family of slices in X_{i+1} – its children – by performing local modifications to the inverse image $\pi_i^{-1}(S) \subset X_{i+1}$. The children of S carry a natural probability measure which treats disjoint local modifications as independent. This

section develops the properties of slices and their children, and then introduces a family of measures $\{\Sigma'_i\}_{i \in \mathbb{Z}}$ on slices.

Let $\{X_i\}$ be an admissible inverse system satisfying Assumption 5.1.

6.1. Slices and their descendants. We recall from Section 2.4 that X_i carries a partial order \preceq .

Definition 6.1. A **partial slice** in X'_i is a finite subset $S \subset V'_i$ which intersects each monotone geodesic $\gamma \subset X_i$ at most once; this is equivalent to saying that no two elements of S are comparable: if $v, v' \in S$ and $v \preceq v'$, then $v = v'$. A **slice** in X'_i is a partial slice which intersects each monotone geodesic precisely once. We denote the set of slices in X'_i and partial slices in X'_i by Slice'_i and PSlice'_i respectively.

The vertex set V'_i is countable, in view of Assumption 5.1. Every partial slice is finite, so this implies that the collection of partial slices is countable. When $i \leq 0$, X'_i is a copy of \mathbb{R} with a standard subdivision, so the slices in X'_i are just singletons $\{v\}$, where $v \in V'_i$.

Note that we cannot have $w \prec x \prec w'$ for $x \in X_i$, $S \in \text{Slice}'_i$, and $w, w' \in S$, because we could concatenate a monotone geodesic segment joining w to w' with monotone rays, obtaining a monotone geodesic which intersects S twice. Therefore we use the notation $x \prec S$ if there is a $w \in S$ such that $x \prec w$. The relations $x \succ S$, $x \preceq S$, and $x \succeq S$ are defined similarly.

A slice $S \in \text{Slice}'_i$ **separates** (respectively **weakly separates**) $x_1, x_2 \in X_i$ if $x_1 \prec S \prec x_2$ or $x_2 \prec S \prec x_1$ (respectively $x_1 \preceq S \preceq x_2$ or $x_2 \preceq S \preceq x_1$).

If $S \in \text{Slice}'_i$, $v \in X_i \setminus S$, then we define

$$\text{Side}(v, S) = \begin{cases} \prec & \text{if } v \prec S \\ \succ & \text{if } v \succ S \end{cases}$$

Slices give rise to monotone sets:

Lemma 6.2. *If $S \in \text{Slice}'_i$, define $S_{\prec} = \{x \in X_i \mid x \preceq S\}$ and $S_{\succ} = \{x \in X_i \mid x \succeq S\}$. Then S_{\prec} and S_{\succ} are both monotone sets with boundary S .*

Proof. Since $S_{\prec}^c = S_{\succ} = \{x \in X_i \mid x \succ S\}$, the monotonicity of S_{\prec} follows immediately from the definition of slices. Similarly for S_{\succ} . \square

Given a vertex $v \in V_i$, we can associate a collection of partial slices in X'_i :

Definition 6.3. If $v \in V_i$, a **child of v** is a maximal partial slice $S' \in \text{PSlice}'_i$ which is contained in the trimmed star $\text{TSt}(v, X_i)$, see Figure 3. In other words, $S' \subset \text{TSt}(v, X_i)$ and precisely one of the following holds:

- (1) $S' = \{v\}$.
- (2) For every vertex $w \in \text{St}(v, X_i)$ with $v \prec w$, S' intersects the edge \overline{vw} in precisely one point, which is an interior point.
- (3) For every vertex $w \in \text{St}(v, X_i)$ with $w \prec v$, S' intersects the edge \overline{vw} in precisely one point, which is an interior point.

We denote the collection of children of v by $\text{Ch}(v)$, and refer to the children in the above cases as **children of type (1), (2), or (3)** respectively.

Note that if $S \in \text{PSlice}'_i$ and $v_1, v_2 \in V_{i+1}$ are distinct vertices lying in $\pi_i^{-1}(S)$, then their trimmed stars are disjoint.

Definition 6.4. If $S \in \text{PSlice}'_i$ is a partial slice, a **child of S** is a subset $S' \subset V'_{i+1}$ obtained by replacing each vertex $v \in \pi_i^{-1}(S) \subset V_{i+1}$ with one of its children, so that S' is a subset of V'_{i+1} . More formally, S' belongs to the image of the “union map”

$$\prod_{v \in \pi_i^{-1}(S)} \text{Ch}(v) \longrightarrow V'_{i+1}$$

which sends $\prod_v (S_v)$ to $\cup_v S_v$. We use $\text{Ch}(S) \subset \text{PSlice}'_{i+1}$ to denote the children of $S \in \text{PSlice}'_i$.

Lemma 6.5. *If S is a partial slice, so is each of its children. Moreover, if $S \in \text{Slice}'_i$ is a slice, so is S' .*

Proof. Suppose S' is a child of the partial slice $S \in \text{PSlice}'_i$, and $\gamma' \subset X_{i+1}$ is a monotone geodesic. Then γ' projects isomorphically to a monotone geodesic $\gamma \subset X_i$, so $\pi_i^{-1}(S) \cap \gamma'$ contains at most one vertex $v' \in V_{i+1}$. From the definition of children, it follows that $\gamma' \cap S'$ contains at most one point. If S is a slice, then S' contains precisely one vertex $v \in V'_{i+1}$, and therefore S' contains a child of $\{v\}$, which will intersect γ' in precisely one point.

□




 Some children of v


FIGURE 3.

Definition 6.6. If $S \in \text{PSlice}'_i$ and $j > i$, then a partial slice $S' \in \text{PSlice}'_j$ is a **descendent of S in X'_j** if there exist $S = S_i, S_{i+1}, \dots, S_j = S'$ such that for all $k \in \{i+1, \dots, j\}$, $S_k \in \text{PSlice}'_k$ and S_k is a child of S_{k-1} ; in other words, S' is an iterated child of S . We denote the collection of such descendents by $\text{Desc}(S, X'_j)$.

Lemma 6.7. For all $i < j$, if $S' \in \text{PSlice}'_j$ is a descendent of $S \in \text{PSlice}'_i$, then $\pi_i(S') \subset \cup_{w \in S} \text{St}^o(w, X'_i)$.

Proof. Suppose $S = S_i, \dots, S_j = S'$, where $S_k \in \text{PSlice}'_k$ and S_{k+1} is a child of S_k for all $i \leq k < j$. Then

$$\pi_k(S_{k+1}) \subset \pi_k \left(\bigcup_{w \in \pi_k^{-1}(S_k)} \text{TSt}(w, X'_{k+1}) \right).$$

Iterating this yields the lemma. \square

6.2. A measure on slices. We now define a measure on Slice'_i for each i , by an iterated diffusion construction. To do this, we first associate with each vertex $v \in V_i$ a probability measure on its children $\text{Ch}(v) \subset \text{PSlice}'_i$.

Definition 6.8. If $v \in V_i$, let $w_{\text{Ch}(v)}$ be the probability measure on $\text{Ch}(v)$ which:

- Assigns measure $\frac{1}{m}$ to the child $\{v\} \in \text{Ch}(v)$ of type (1).
- Uniformly distributes measure $\frac{1}{2} \cdot \frac{(m-1)}{m}$ among the children of type (2). Equivalently, for each vertex $\hat{v} \in V_i$ adjacent to v with $v \prec \hat{v}$, we take the uniform measure on the $(m-1)$ vertices in V'_i which are interior points of the edge $\overline{v\hat{v}}$, take the product of these measures as \hat{v} ranges over

$$\{\bar{v} \in V_i \mid \bar{v} \in \text{St}(v, X_i), v \prec \bar{v}\}$$

and then multiply the result by $\frac{1}{2} \cdot \frac{(m-1)}{m}$.

- Uniformly distributes measure $\frac{1}{2} \cdot \frac{(m-1)}{m}$ among the children of type (3).

Note that if $v' \in V'_i$ belongs to the trimmed star of v , then the $w_{\text{Ch}(v)}$ measure of the children of v which contain v' is

$$(6.9) \quad w_{\text{Ch}(v)}(\{S \in \text{Ch}(v) \mid v' \in S\}) = \begin{cases} \frac{1}{m} & \text{if } v' = v \\ \frac{1}{2m} & \text{if } v' \neq v \end{cases}$$

Using the measures $w_{\text{Ch}(v)}$ we define a measure on the children of a slice:

Definition 6.10. If $S \in \text{Slice}'_i$, we define a probability measure K_S on $\text{Ch}(S)$ as follows. We take the product measure $\prod_{v \in \pi_i^{-1}(S)} w_{\text{Ch}(v)}$ on $\prod_{v \in \pi_i^{-1}(S)} \text{Ch}(v)$, and push it forward under the union map

$$\prod_{v \in \pi_i^{-1}(S)} \text{Ch}(v) \rightarrow \text{Slice}'_{i+1}.$$

In probabilistic language, for each $v \in \pi_i^{-1}(S)$, we independently choose a child of v according to the distribution $w_{\text{Ch}(v)}$, and then take the union of the resulting children. Note that this is well-defined because the inverse image of any slice is nonempty.

Now given a measure ν on Slice'_i , we diffuse it to a measure ν' on Slice'_{i+1} :

$$(6.11) \quad \nu' = \sum_{S \in \text{Slice}'_i} K_S \nu(S).$$

If we view the collection $\{K_S\}_{S \in \text{Slice}'_i}$ as defining a kernel

$$K_i : \text{Slice}'_i \times \text{Slice}'_{i+1} \rightarrow [0, 1]$$

by the formula $K_i(S, S') = K_S(S')$, then the associated diffusion operator K_i is given by

$$(6.12) \quad K_i(\nu)(S') = \sum_{S \in \text{Slice}'_i} K(S, S') \nu(S).$$

When $i < 0$, then this sum will be finite for any measure ν since $K(S, S') \neq 0$ for only finitely many S .

Lemma 6.13. *When $i \geq 0$, the sum will be finite provided ν is supported on the descendants of slices in X'_0 .*

Proof. For a given $S' \in \text{Slice}'_{i+1}$, the summand $K(S, S')\nu(S)$ is nonzero only if S is a descendent of a slice $\{v\} \in V'_0$ and S' is a child of S . By Lemma 6.7 this means that $\pi_0(S') \subset \text{St}^o(v, X'_0)$, so there are only finitely many possibilities for such S . \square

Definition 6.14. For $i \leq 0$, let Σ'_i the measure on Slice'_i which assigns measure $m^{-(i+1)}$ to each slice in $\text{Slice}'_i = V'_i$. For $i > 0$ we define a measure Σ'_i on Slice'_i inductively by $\Sigma'_i = K_{i-1}(\Sigma'_{i-1})$. This is well-defined by Lemma 6.13.

For every $S \in \text{Slice}'_i$ and every $j > i$, we may also obtain a well-defined probability measure on Slice'_j which is supported on the descendants of S , by the formula

$$(6.15) \quad K_{j-1} \circ \dots \circ K_i(\delta_S),$$

where δ_S is a Dirac mass on S . Using this probability measure, we may speak of the measure of descendants of $S \in \text{Slice}'_j$.

7. ESTIMATES ON THE FAMILY OF MEASURES $\{\Sigma'_i\}_{i \in \mathbb{Z}}$

In this section we will prove (mostly by induction arguments) several estimates on the slice/cut measures $\{\Sigma'_i\}$ and cut metrics $\{d_{\Sigma'_i}\}$ that will be needed in Section 8.

We first observe that the slices passing through a vertex $v \in V'_i$ have measure $m^{-(i+1)}$:

Lemma 7.1. *For all $i \in \mathbb{Z}$, and every $v \in V'_i$, the Σ'_i -measure of the collection of slices containing v is precisely $m^{-(i+1)}$:*

$$\Sigma'_i(\{S \in \text{Slice}'_i \mid v \in S\}) = m^{-(i+1)}.$$

Proof. When $i \leq 0$ this reduces to the definition of Σ'_i . So pick $i > 0$, $v \in V'_i$, and assume inductively that the lemma is true for $i - 1$.

Case 1. $v \in V_i$. In this case, if a slice $S \in \text{Slice}'_{i-1}$ has a child $S' \in \text{Slice}'_i$ containing v , then $\pi_{i-1}(v) \in S$. By Definition 6.8, for such an S , the fraction of its children containing v is precisely $\frac{1}{m}$. Therefore by the induction hypothesis we have

$$\Sigma'_i(\{S' \mid v \in S'\}) = \frac{1}{m} \Sigma'_{i-1}(\{S \in \text{Slice}'_{i-1} \mid \pi_{i-1}(v) \in S\}) = m^{-(i+1)}.$$

Case 2. $v \notin V_i$. Then v belongs to a unique edge $\overline{w_1 w_2} \subset X_i$, where $w_1, w_2 \in V_i$. In this case, a slice $S \in \text{Slice}'_{i-1}$ has a child $S' \in \text{Slice}'_i$ containing v if and only if S contains $\pi_{i-1}(w_1)$ or $\pi_{i-1}(w_2)$. Since these possibilities are mutually exclusive (from the definition of slice), and each contributes a measure $\frac{1}{2}m^{-(i+1)}$ by the induction hypothesis and Definition 6.8, the lemma follows. \square

Recall that by Lemma 6.2, for every $S \in \text{Slice}'_i$ the subset $S_{\leq} = \{x \in X_i \mid x \leq S\}$ is a monotone subset of X_i .

Definition 7.2. Viewing Σ'_i as a cut measure on X_i via the identification $S \longleftrightarrow S_{\leq}$, we let $d_{\Sigma'_i}$ denote the corresponding cut metric on X_i . Equivalently, for $x_1, x_2 \in X_i$,

$$d_{\Sigma'_i}(x_1, x_2) = \sum_{S \in \text{Slice}'_i} d_{S_{\leq}}(x_1, x_2) \Sigma'_i(S)$$

where

$$d_{S_{\leq}}(x_1, x_2) = |\chi_{S_{\leq}}(x_1) - \chi_{S_{\leq}}(x_2)|$$

$$= \begin{cases} 1 & \text{if } x_1 \preceq S \prec x_2 \text{ or } x_2 \preceq S \prec x_1 \\ 0 & \text{otherwise} \end{cases}$$

Lemma 7.3. *If $x'_1, x'_2 \in X_{i+1}$, and $\pi_i(x'_j) = x_j \in X_i$, then*

$$(7.4) \quad |d_{\Sigma'_{i+1}}(x'_1, x'_2) - d_{\Sigma'_i}(x_1, x_2)| \leq 4m^{-(i+1)}.$$

Proof. For $j \in \{1, 2\}$ let S_j be the collection of slices $S \in \text{Slice}'_i$ which have a child $S' \in \text{Slice}'_{i+1}$ such that $\text{Side}(x_j, S) \neq \text{Side}(x'_j, S')$. From the definition of children, it follows that if $S \in S_j$, then x_j lies in $\text{TSt}(v, X'_i)$ for some $v \in S$. Thus, if $\overline{w_1 w_2}$ is an edge of X'_i containing x_j , then $S_j \subset \{S \in \text{Slice}'_i \mid S \cap \{w_1, w_2\} \neq \emptyset\}$. By Lemma 7.1, we have $\Sigma'_i(S_j) \leq 2m^{-(i+1)}$. Now by the definition of the cut metrics, we get

$$|d_{\Sigma'_{i+1}}(x'_1, x'_2) - d_{\Sigma'_i}(x_1, x_2)| \leq \Sigma'_i(S_1 \cup S_2) \leq 4m^{-(i+1)}.$$

□

Lemma 7.5 (Persistence of sides). *Suppose $x \in X_i$, $S \in \text{Slice}'_i$, and $x \notin \cup_{v \in S} \text{St}^o(v, X'_i)$. Then for every $j \geq i$, every $x' \in X_j$ with $\pi_i(x') = x$, and for every descendent $S' \in \text{Slice}'_j$ of S , we have*

$$x' \notin \cup_{v \in S'} \text{St}^o(v, X'_j) \quad \text{and} \quad \text{Side}(x', S') = \text{Side}(x, S).$$

Proof. First suppose $j = i + 1$. Then $x \notin \cup_{v \in \pi_i^{-1}(S)} \text{St}^o(v, X_{i+1})$ because $\pi_i : X_{i+1} \rightarrow X'_i$ is a simplicial mapping. Clearly this implies $\text{Side}(x', S') = \text{Side}(x', \pi_{i+1}^{-1}(S)) = \text{Side}(x, S)$.

The $j > i + 1$ case now follows by induction.

□

Lemma 7.6 (Persistence of separation). *There is a constant $A = A(m) \in (0, 1)$ with the following property. Suppose $j > i$, $x_1, x_2 \in X_i$, $x'_1, x'_2 \in X_j$, and $\pi_i(x'_1) = x_1$, $\pi_i(x'_2) = x_2$. Suppose in addition that*

- $S \in \text{Slice}'_i$ is a slice which weakly separates x_1 and x_2 .
- $x_2 \notin \cup_{v \in S} \text{St}^o(v, X'_i)$.

Then the measure of the collection of descendants $S' \in \text{Desc}(S, X'_j)$ which separate x'_1 and x'_2 is at least A ; here we refer to the probability measure on $\text{Desc}(S, X_j)$ that was defined in equation (6.15).

Proof. Since S weakly separates x_1 and x_2 but $x_2 \notin S$, without loss of generality we may assume that $x_1 \preceq S \prec x_2$, since the case $x_2 \prec S \preceq x_1$ is similar.

If $x_1 \notin \cup_{v \in S} \text{St}^o(v, X'_i)$, then by Lemma 7.5, for all $S' \in \text{Desc}(S, X'_j)$ we have $x'_1 \prec S' \prec x'_2$, so we are done in this case.

Therefore we assume that there exist $v \in S$ and $v' \in \pi_i^{-1}(v)$ such that $x_1 \in \text{St}^o(v, X'_i)$, $x'_1 \in \text{St}(v', X_{i+1})$, and $x'_1 \preceq v'$. If $S' \in \text{Slice}'_{i+1}$ is a child of S containing a child of v' of type (2), then $x'_1 \notin \cup_{w \in S'} \text{St}^o(w, X'_{i+1})$; moreover the collection of such slices S' form a fraction at least $\frac{1}{2} \frac{m-1}{m}$ of the children of S . Applying Lemma 7.5 to each such slice S' , we conclude that for every $S'' \in \text{Desc}(S', X'_j)$, we have $x'_1 \prec S'' \prec x'_2$. This proves the lemma. \square

Lemma 7.7. *Suppose $x_1, x_2 \in X_i$, $j > i$, $x'_1, x'_2 \in X_j$, $\pi_i(x'_1) = x_1$, $\pi_i(x'_2) = x_2$, and $\{x_1, x_2\}$ is not contained in the trimmed star of any vertex $v \in V_i$. Then $d_{\Sigma'_j}(x'_1, x'_2) \geq A m^{-(i+2)}$, where A is the constant from Lemma 7.6.*

Proof. Choose $v_1 \in V_i$ such that $x_1 \in \text{TSt}(v_1, X_i)$.

Observe that of the children W of v_1 , a measure at least $\frac{1}{2m}$ lie weakly on each side of x_1 and satisfy $x_2 \notin \cup_{v \in W} \text{St}^o(v, X'_i)$, in view of our assumption on x_1 and x_2 , i.e.

$$w_{\text{Ch}(v_1)}(\{W \in \text{Ch}(v_1) \mid W \preceq x_1, x_2 \notin \cup_{v \in W} \text{St}^o(v, X'_i)\}) \geq \frac{1}{2m},$$

$$w_{\text{Ch}(v_1)}(\{W \in \text{Ch}(v_1) \mid W \succeq x_1, x_2 \notin \cup_{v \in W} \text{St}^o(v, X'_i)\}) \geq \frac{1}{2m}.$$

Suppose $S \in \text{Slice}'_{i-1}$ and $v_1 \in \pi_{i-1}^{-1}(S)$. Then each child $S' \in \text{Slice}'_i$ of S contains some child of v_1 , and $\text{Side}(x_2, S')$ is independent of this choice, because x_2 lies outside $\text{TSt}(v_1, X_i)$. Furthermore, if $x_2 \in \text{St}(v_2, X_i)$ for some $v_2 \in (V_i \cap \pi_{i-1}^{-1}(S)) \setminus \{v_1\}$, then S' contains a child of v_2 , and a fraction at least $\frac{1}{m}$ of this set of children $W \in \text{Ch}(v_2)$ satisfies $x_2 \notin \cup_{v \in W} \text{St}^o(v, X'_i)$. Thus a fraction at least $\frac{1}{2m^2}$ of the children of S satisfy the assumptions of Lemma 7.6.

Since the set of $S \in \text{Slice}'_{i-1}$ with $v_1 \in \pi_{i-1}^{-1}(S)$ has Σ'_{i-1} -measure m^{-i} by Lemma 7.1, by the preceding reasoning, we conclude that $d_{\Sigma'_j}(x'_1, x'_2) \geq A m^{-(i+2)}$. \square

Lemma 7.8. *Suppose $i, j \in \mathbb{Z}$, $i \leq j$, $x_1, x_2 \in X_j$, e is an edge of X_i , and $\pi_i(x_1), \pi_i(x_2) \in e$. Then*

$$(7.9) \quad d_{\Sigma'_j}(x_1, x_2) \leq m^{-i}.$$

Proof. Let v_1, v_2 be the endpoints of e , where $v_1 \prec v_2$. We may assume without loss of generality that $\pi_i(x_1) \preceq \pi_i(x_2)$.

By Definition 7.2, the distance $d_{\Sigma'_j}(x_1, x_2)$ is the Σ'_j -measure of the set

$$Y = \{S \in \text{Slice}'_j \mid x_1 \preceq S \prec x_2 \text{ or } x_2 \preceq S \prec x_1\}.$$

If $S \in \text{Slice}'_i$ and $S \cap e = \emptyset$, then S does not weakly separate $\pi_i(x_1), \pi_i(x_2)$, so by Lemma 7.5, no descendent $S' \in \text{Desc}(S, X'_j)$ can weakly separate x_1 and x_2 , i.e. $Y \cap \text{Desc}(S, X'_j) = \emptyset$. For $v \in V'_i$ let

$$\text{Slice}'_i(v) = \{S \in \text{Slice}'_i \mid v \in S\}$$

and let $\Sigma'_i \llcorner \text{Slice}'_i(v)$ be the restriction of Σ'_i to $\text{Slice}'_i(v)$. Thus, using the diffusion operators K_l from (6.11), the above observation implies that

$$\begin{aligned} \Sigma'_j(Y) &= K_{j-1} \circ \dots \circ K_i(\Sigma'_i(Y)) \\ &= \sum_{v \in e \cap V'_i} K_{j-1} \circ \dots \circ K_i(\Sigma'_i \llcorner \text{Slice}'_i(v))(Y) \end{aligned} \quad (7.10)$$

$$\leq (m-1)m^{-(i+1)} + \sum_{v \in \{v_1, v_2\}} K_{j-1} \circ \dots \circ K_i(\Sigma'_i \llcorner \text{Slice}'_i(v))(Y),$$

because the mass of $\Sigma'_i \llcorner \text{Slice}'_i(v)$ is $m^{-(i+1)}$ by Lemma 7.1. The remainder of the proof is devoting to showing that the total contribution from the last two terms in (7.10) is at most $m^{-(i+1)}$.

Suppose $\pi_i(x_1) \notin \text{St}^o(v_1, X'_i)$. If $S \in \text{Slice}'_i(v_1)$, then no descendent $S' \in \text{Desc}(S, X'_j)$ can weakly separate x_1, x_2 by Lemma 7.5. Therefore

$$K_{j-1} \circ \dots \circ K_i(\Sigma'_i \llcorner \text{Slice}'_i(v_1))(Y) = 0,$$

so we are done in this case. Hence we may assume that $\pi_i(x_1) \in \text{St}^o(v_1, X'_i)$, and by similar reasoning, that $\pi_i(x_2) \in \text{St}^o(v_2, X'_i)$. This implies by Lemma 7.5 that if $v_1 \in S \in \text{Slice}'_i$ and $S' \in \text{Desc}(S, X'_j)$ then $S' \prec x_2$; similarly, if $v_2 \in S \in \text{Slice}'_i$ and $S' \in \text{Desc}(S, X'_j)$, then $x_1 \prec S'$.

By (7.10), the lemma now reduces to the following two claims:

Claim 1. $K_{j-1} \circ \dots \circ K_i(\Sigma'_i \sqcup \text{Slice}'_i(v_1))(Y) \leq \frac{1}{2}(m^{-(i+1)} + m^{-(j+1)}).$

Claim 2. $K_{j-1} \circ \dots \circ K_i(\Sigma'_i \sqcup \text{Slice}'_i(v_2))(Y) \leq \frac{1}{2}(m^{-(i+1)} - m^{-(j+1)}).$

Proof of Claim 1. For each $i \leq k \leq j$, let w'_k be the unique vertex in V'_k such that $\pi_k(x_1) \in \text{St}^o(w'_k, X'_k)$ and $w'_k \preceq \pi_k(x_1)$. Likewise, let w_k be the unique vertex in V_k such that $\pi_k(x_1) \in \text{St}^o(w_k, X_k)$ and $w_k \preceq \pi_k(x_1)$. Thus $w'_k \in \text{St}^o(w_k, X_k)$. Let k_0 be the maximum of the integers $k \in [i, j]$ such that $\pi_i(w'_k) = v_1$ for all $i \leq k \leq k_0$. It follows that $w'_k = w_k$ for all $i \leq k \leq k_0$.

For $i \leq k \leq j$, let A_k be the collection of slices $S \in \text{Slice}'_k$ which contain w_k , and let B_k be the collection of slices $S \in \text{Slice}'_k$ which contain a child of w_k of type (3). We now define a sequence of measures $\{\alpha_k\}_{i \leq k \leq k_0}$ inductively as follows. Let α_i be the restriction of Σ'_i to $\{S \in \text{Slice}'_i \mid v_1 \in S\}$. For $k < k_0$, we define α_{k+1} to the restriction of $K_k \alpha_k$ to $\text{Slice}'_{k+1} \setminus B_{k+1}$, where K_k is the diffusion operator (6.11). Since by Lemma 7.5 the set of descendants $\text{Desc}(B_k, X'_j)$ is disjoint from Y , it follows that

$$(7.11) \quad K_{j-1} \circ \dots \circ K_i(\Sigma'_i \sqcup \text{Slice}'_i(v_1)) \sqcup Y = (K_{j-1} \circ \dots \circ K_k \alpha_k) \sqcup Y$$

for all $k \leq k_0$.

Note that by the definition of the diffusion operator K_{k-1} we have

$$(7.12) \quad \alpha_k(A_k) \geq \frac{1}{m} \cdot \alpha_{k-1}(A_{k-1})$$

$$(7.13) \quad (K_{k-1} \alpha_{k-1})(B_k) \geq \frac{(m-1)}{2m} \cdot \alpha_{k-1}(A_{k-1})$$

for all $i < k \leq k_0$. This yields $\alpha_k(A_k) \geq m^{-(k+1)}$ for all $i \leq k \leq k_0$. Hence for all $i < k \leq k_0$ we get

$$\begin{aligned} \alpha_k(\text{Slice}'_k) &= \alpha_{k-1}(\text{Slice}'_{k-1}) - (K_{k-1} \alpha_{k-1})(B_k) \\ &\leq \alpha_{k-1}(\text{Slice}'_{k-1}) - \frac{(m-1)}{2m} \cdot m^{-k} \end{aligned}$$

and so

$$\begin{aligned} \alpha_{k_0}(\text{Slice}'_{k_0}) &\leq m^{-(i+1)} - \frac{(m-1)}{2m} \cdot (m^{-(i+1)} + \dots + m^{-k_0}) \\ &= \frac{1}{2} m^{-(i+1)} + \frac{1}{2} m^{-(k_0+1)}. \end{aligned}$$

This gives Claim 1 when $k_0 = j$, by (7.11).

We now assume that $k_0 < j$. Then $w_{k_0+1} \prec w'_{k_0+1}$. By Lemma 7.5, every descendent $S' \in \text{Slice}'_j$ of a slice $S \in A_{k_0+1} \cup B_{k_0+1}$ will satisfy

$S' \prec x_1$, so $S' \notin Y$. Therefore if we define α_{k_0+1} to be the restriction of $K_{k_0}\alpha_{k_0}$ to $\text{Slice}'_{k_0+1} \setminus (A_{k_0+1} \cup B_{k_0+1})$, then

$$K_{j-1} \circ \dots \circ K_i(\Sigma'_i \sqcup \text{Slice}'_i(v_1))(Y) = K_{j-1} \circ \dots \circ K_{k_0+1}(\alpha_{k_0+1})(Y).$$

Also, as in (7.12)–(7.13), we get

$$\begin{aligned} K_{k_0}\alpha_{k_0}(A_{k_0+1} \sqcup B_{k_0+1}) &\geq \left(\frac{1}{m} + \frac{(m-1)}{2m}\right) \alpha_{k_0}(A_{k_0}) \\ &\geq \left(\frac{1}{m} + \frac{(m-1)}{2m}\right) m^{-(k_0+1)}. \end{aligned}$$

Therefore

$$\begin{aligned} K_{j-1} \circ \dots \circ K_{k_0+1}(\alpha_{k_0+1})(Y) &\leq \alpha_{k_0+1}(\text{Slice}'_{k_0+1}) \\ &= \alpha_{k_0}(\text{Slice}'_{k_0}) - K_{k_0}\alpha_{k_0}(A_{k_0+1} \sqcup B_{k_0+1}) \\ &\leq \frac{1}{2}m^{-(i+1)} + \frac{1}{2}m^{-(k_0+1)} - \left(\frac{1}{m} + \frac{(m-1)}{2m}\right) m^{-(k_0+1)} \\ &\leq \frac{1}{2}m^{-(i+1)} - \frac{1}{2}m^{-(k_0+1)} \end{aligned}$$

so Claim 1 holds.

Proof of Claim 2. The proof is similar to that of Claim 1, except that one replaces v_1 with v_2 , and reverses the orderings. However, in the case when $k_0 = j$, one simply notes that any slice $S' \in A_{k_0} = A_j$ satisfies $x_1 \preceq S'$, $x_2 \preceq S'$, so $S' \notin Y$. Therefore we may remove the measure contributed by A_{k_0} from our estimate, making it smaller by $m^{-(j+1)}$.

□

Corollary 7.14. *Suppose $i, j \in \mathbb{Z}$, $i \leq j$, $x_1, x_2 \in X_j$, $v \in V_i$, and $\{x_1, x_2\} \subset (\pi_i^j)^{-1}(\text{St}(v, X_i))$. Then*

$$(7.15) \quad d_{\Sigma_j'}(x_1, x_2) \leq 2m^{-i}.$$

Proof. First suppose there is an $x \in X_j$ such that $\pi_i^j(x) = v$. Then $\{\pi_i^j(x), \pi_i^j(x_1)\}$ lies in an edge of X_i , so by Lemma 7.8 we have $d_{\Sigma_j'}(x, x_1) \leq m^{-i}$, and similarly $d_{\Sigma_j'}(x, x_2) \leq m^{-i}$. Therefore (7.15) holds.

In general, construct a new admissible inverse system $\{Y_k\}$ satisfying Assumption 5.1 by letting Y_k be the disjoint union of X_k with a copy of \mathbb{R} when $i < k \leq j$, and $Y_k = X_k$ otherwise. Then extend the projection map $\pi_i : X_{i+1} \rightarrow X_i$ to $\pi_i : Y_{i+1} \rightarrow Y_i = X_i$ by mapping $Y_{i+1} \setminus X_{i+1} \simeq \mathbb{R}$

to a monotone geodesic containing v . Then for $i < k < j$ extend $\pi_j : X_{j+1} \rightarrow X_j$ to $\pi_j : Y_{j+1} \rightarrow Y_j$ by mapping $Y_{j+1} \setminus X_{j+1} \simeq \mathbb{R}$ isomorphically to $Y_j \setminus X_j \simeq \mathbb{R}$. Then there is a system of measures $\{\Sigma'_{k,Y}\}$ for the inverse system $\{Y_k\}$, and it follows that the associated cut metric $d_{\Sigma'_j}^Y$ is the same for pairs $x_1, x_2 \subset X_j \subset Y_j$. Since v belongs to the image of $\pi_i^j : Y_j \rightarrow Y_i$, we have $d_{\Sigma'_j}^X(x_1, x_2) = d_{\Sigma'_j}^Y(x_1, x_2) \leq 2m^{-i}$. \square

8. PROOF OF THEOREM 1.16 UNDER ASSUMPTION 5.1

We will define a sequence $\{\rho_i : X_\infty \times X_\infty \rightarrow [0, \infty)\}$ of pseudo-distances on X_∞ , such that ρ_i is induced by a map $X_\infty \rightarrow L_1$, and ρ_i converges uniformly to some $\rho_\infty : X_\infty \times X_\infty \rightarrow [0, \infty)$. By a standard argument, this yields an isometric embedding $(X_\infty, \rho_\infty) \rightarrow L_1$. (The theory of ultralimits [HM82, BL00] implies the metric space (X_∞, ρ_∞) isometrically embeds in an ultralimit V of L_1 spaces; by Kakutani's theorem [Kak39] the space V is isometric to an L_1 space, and so ρ_∞ isometrically embeds in L_1 .) To complete the proof, it suffices to verify that ρ_∞ has the properties asserted by the theorem. (Alternately, one may construct a cut measure Σ_∞ on X_∞ as weak limit, and use the corresponding cut metric to provide the embedding to L_1 .)

Let $\rho_i = (\pi_i^\infty)^* d_{\Sigma'_i}$ be the pullback of $d_{\Sigma'_i}$ to X_∞ . By Lemma 7.3 we have $|\rho_{i+1} - \rho_i| \leq 4m^{-(i+1)}$, so the sequence $\{\rho_i\}$ converges uniformly to a pseudo-distance $\rho_\infty : X_\infty \times X_\infty \rightarrow [0, \infty)$.

Lemma 8.1.

- (1) $\rho_\infty(x, x') \leq \bar{d}_\infty(x, x')$.
- (2) $\rho_\infty(x, x') \geq \frac{A}{2m^3} \cdot \bar{d}_\infty(x, x')$.
- (3) $\rho_\infty(x, x') = \bar{d}_\infty(x, x')$ if x, x' lie on a monotone geodesic.

Proof. (1). Suppose $x, x' \in X_\infty$, and for some $i \in \mathbb{Z}$ the projections $\{\pi_i(x), \pi_i(x')\}$ are contained in $\text{St}(v, X_i)$. By Corollary 7.14 we have $\rho_\infty(x, x') \leq 2m^{-i}$. Now (1) follows from the definition of \bar{d}_∞ .

(2). Suppose $x \neq x'$, and let $j \in \mathbb{Z}$ be the minimum of the indices $k \in \mathbb{Z}$ such that $\pi_k(x), \pi_k(x')$ are not contained in the trimmed star of any vertex $v \in X_k$. Then $\bar{d}_\infty(x, x') \leq 2m^{-(j-1)}$ by Lemma 2.6, while $\rho_\infty(x, x') \geq Am^{-(j+2)}$ by Lemma 7.7. Thus

$$\rho_\infty(x, x') \geq \frac{A}{2m^3} \cdot \bar{d}_\infty(x, x').$$

(3). If $x, x' \in X_\infty$ lie on a monotone geodesic γ , then γ will project homeomorphically under π_i to a monotone geodesic $\pi_i(\gamma)$, which contains at least $\bar{d}_\infty(x_1, x_2)m^{(i+1)} - 2$ vertices of V'_i . By Lemma 7.1 we have $\rho_i(x_1, x_2) \geq \bar{d}_\infty(x_1, x_2) - 2m^{-(i+1)}$. Since i was arbitrary we get $\rho_\infty(x, x') = \bar{d}_\infty(x, x')$. \square

This completes the proof of Theorem 1.16 under Assumption 5.1.

9. THE PROOF OF THEOREM 1.16, GENERAL CASE

We recall the three conditions from Assumption 5.1:

- (1) π_i is finite-to-one for all $i \in \mathbb{Z}$.
- (2) $X_i \simeq \mathbb{R}$ and $\pi_i : X_{i+1} \rightarrow X'_i$ is an isomorphism for all $i \leq 0$.
- (3) For every $i \in \mathbb{Z}$, every vertex $v \in V_i$ has neighbors $v_\pm \in V_i$ with $v_- \prec v \prec v_+$. Equivalently, X_i is a union of monotone geodesics (see Definition 2.12).

In this section these three conditions will be removed in turn.

9.1. Removing the finiteness assumption. We now assume that $\{X_i\}$ is an admissible inverse system satisfying conditions (2) and (3) of Assumption 5.1, but not necessarily the finiteness condition (1).

To prove Theorem 1.16 without the finiteness assumption, we observe that the construction of the distance function $\rho_\infty : X_\infty \times X_\infty \rightarrow \mathbb{R}$ can be reduced to the case already treated, in the sense that for any two points $x_1, x_2 \in X_\infty$, we can apply the construction of the cut metrics to finite valence subsystems, and the resulting distance $\rho_\infty(x_1, x_2)$ is independent of the choice of subsystem. The proof is then completed by invoking the main result of [DCK72]. We now give the details.

Definition 9.1. A **finite subsystem** of the inverse system $\{X_j\}$ is a collection of subcomplexes $\{Y_j \subset X_j\}_{j=-\infty}^i$, for some $i \geq 0$, such that $\pi_j(Y_{j+1}) \subset Y_j$ for all $j < i$, and $\{Y_j\}_{j \leq i}$ satisfies Assumption 5.1 for indices $\leq i$. In other words:

- (1) π_j is finite-to-one for all $j \leq i$.
- (2) $Y_j \simeq \mathbb{R}$ and $\pi_j : Y_{j+1} \rightarrow Y'_j$ is an isomorphism for all $j \leq 0$.
- (3) Y_j is a union of (complete) monotone geodesics for all $j \leq i$.

Suppose $i \geq 0$ and V is a finite subset of $V'_i \subset X_i$. Then there exists a finite subsystem $\{Y_j\}_{j=-\infty}^i$ such that Y_i contains V . One may obtain such a system by letting Y_i be a finite union of monotone geodesics in X_i which contains V , and taking $Y_j = \pi_j^i(Y_i)$ for $j < i$. The inductive construction of the slice measures in the finite valence case may be applied to the finite subsystem $\{Y_j\}$, to obtain a sequence of slice measures which we denote by $\Sigma'_{j,Y}$, to emphasize the potential dependence on Y .

Lemma 9.2. *Suppose $i \geq 0$, $V \subset V'_i$ is a finite subset, and let $\{Y_j\}_{j \leq i_Y}$ and $\{Z_j\}_{j \leq i_Z}$ be finite subsystems of $\{X_j\}$, where $i \leq \min(i_Y, i_Z)$ and $V \subset Y_i \cap Z_i$. If $\Sigma'_{i,Y}$, $\Sigma'_{i,Z}$ denote the respective slice measures, then*

$$\Sigma'_{i,Y}(\{S \in \text{Slice}'_{i,Y} \mid S \supset V\}) = \Sigma'_{i,Z}(\{S \in \text{Slice}'_{i,Z} \mid S \supset V\}),$$

i.e. the slice measure does not depend on the choice of subsystem containing V .

Proof. If $i \leq 0$ then $Y_i = Z_i$ and $\Sigma'_{i,Y} = \Sigma'_{i,Z}$ by construction. So assume that $i > 0$, and that the lemma holds for all finite subsets of $V'_{\bar{i}}$ for all $\bar{i} < i$.

Suppose $S' \in \text{Slice}'_{i,Y}$ is a child of $S \in \text{Slice}'_{i-1,Y}$, and $V \subset S'$. Then for every $v \in V$, either $v \in V_i$, in which case $\pi_{i-1}(v) \in S$, or $v \in V'_i \setminus V_i$, in which case v is an interior point of some edge $\overline{u_1 u_2}$ of $Y_i \subset X_i$, and S contains precisely one of the points $\pi_{i-1}(u_1)$, $\pi_{i-1}(u_2)$. By the definition of $\Sigma'_{i,Y}$ given by (6.11):

$$\begin{aligned} & \Sigma'_{i,Y}(\{S' \in \text{Slice}'_{i,Y} \mid S' \supset V\}) \\ (9.3) \quad &= \sum_{S \in \text{Slice}'_{i-1,Y}} K_S(\{S' \in \text{Slice}'_{i,Y} \mid S' \supset V\}) \Sigma'_{i-1,Y}(S) \end{aligned}$$

By the above observation, the nonzero terms in the sum come from the slices $S \in \text{Slice}'_{i-1,Y}$ which contain precisely one of a finite collection of finite subsets $\bar{V}_1, \dots, \bar{V}_k \subset V'_{i-1}$. If $S \in \text{Slice}'_{i-1,Y}$ contains \bar{V}_l , then from the definition of K_S , the quantity $K_S(\{S' \in \text{Slice}'_{i,Y} \mid S' \supset V\})$ depends only on V_l . Therefore by the induction assumption, it follows that the nonzero terms in (9.3) will be the same as the corresponding terms in the sum defining $\Sigma'_{i,Z}(\{S' \in \text{Slice}'_{i,Y} \mid S' \supset V\})$.

□

Lemma 9.4. *If $\{Y_j\}_{j \leq i}$ is a finite subsystem such that Y_i contains $\{x_1, x_2\} \subset X_i$, then the cut metric $d_{\Sigma'_{i,Y}}(x_1, x_2)$ does not depend on the choice of $\{Y_j\}_{j \leq i}$.*

Proof. Let $\gamma_1, \gamma_2 \subset Y_i$ be monotone geodesics containing x_1 and x_2 respectively. Then $d_{\Sigma'_{i,Y}}(x_1, x_2)$ is the total $\Sigma'_{i,Y}$ -measure of the slices $S \in \text{Slice}'_{i,Y}$ such that either $x_1 \preceq S \prec x_2$ or $x_2 \preceq S \prec x_1$. But every such slice S contains precisely one point from γ_1 , and one point from γ_2 . As the choice of γ_1, γ_2 was arbitrary, Lemma 9.2 implies that cut metric $d_{\Sigma'_{i,Y}}(x_1, x_2)$ does not change when we pass from $\{Y_j\}_{j \leq i}$ to another subsystem which contains $\{Y_j\}_{j \leq i}$. This implies the lemma, since the union of any two finite subsystems containing $\{x_1, x_2\}$ is a finite subsystem which assigns the same cut metric to (x_1, x_2) . \square

We now define a sequence of pseudo-distances $\{\rho_i : X_\infty \times X_\infty \rightarrow [0, \infty)\}$ by letting $\rho_i(x_1, x_2) = d_{\Sigma_{i,Y}}(\pi_i^\infty(x_1), \pi_i^\infty(x_2))$ where $\{Y_j\}_{j \leq i}$ is any finite subsystem containing $\{\pi_i^\infty(x_1), \pi_i^\infty(x_2)\}$. By Lemma 9.4 the pseudo-distance is well-defined. As in the finite valence case:

- Lemma 7.3 implies that $\{\rho_i\}$ converges uniformly to a pseudo-distance $\rho_\infty : X_\infty \times X_\infty \rightarrow [0, \infty)$.
- $\frac{A}{2m^3} \bar{d}_\infty \leq \rho_\infty \leq \bar{d}_\infty$, since this may be verified for each pair of points $x_1, x_2 \in X_\infty$ at a time, by using finite subsystems.
- If $V \subset X_\infty$ is a finite subset, then the restriction of ρ_i to V embeds isometrically in L_1 for all i , and hence the same is true for ρ_∞ .

By the main result of [DCK72], if Z is a metric space such that every finite subset isometrically embeds in L_1 , then Z itself isometrically embeds in L_1 . Therefore (X_∞, ρ_∞) isometrically embeds in L_1 .

9.2. Removing Assumption 5.1(2). Now suppose $\{X_i\}$ is an admissible inverse system satisfying Assumption 5.1(3), i.e. it is a union of monotone geodesics. We will reduce to the case treated in Section 9.1 by working with balls, and then take an ultralimit as the radius tends to infinity.

Lemma 9.5. *Suppose $p \in X_\infty$, $R \in (0, \infty)$ and $R < m^{-(i+1)}$. Then there is an admissible inverse system $\{Z_j\}$ satisfying (2) and (3) of Assumption 5.1, and an isometric embedding of the rescaled ball:*

$$\phi : (B(p, R), m^{(i-1)} \bar{d}_\infty) \rightarrow Z_\infty$$

which preserves the partial order, i.e. if $x, y \in B(p, R)$ and $x \preceq y$, then $\phi(x) \preceq \phi(y)$.

Proof. Since $R < m^{-(i+1)}$, by Lemma 2.4 there is a $v \in V_i$ such that $\pi_i(B(p, R)) \subset \text{TSt}(v, X_i)$.

We now construct an inverse system $\{Y_j\}$ as follows. For $j \geq i$, we let Y_j be the inverse image of $\text{St}(v, X_i)$ under the projection $\pi_i^j : X_j \rightarrow X_i$. We let $Y_j \simeq \mathbb{R}$ for $j < i$. To define the projection maps, we take $\pi_j^Y = \pi_i^X|_{Y_{j+1}}$ for $j \geq i$, and let $\pi_j^Y : Y_{j+1} \rightarrow Y_j'$ be a simplicial isomorphism for $j < i - 1$. Finally, we take $\pi_{i-1}^Y : Y_i = \text{St}(v, X_j) \rightarrow Y_{i-1}' \simeq \mathbb{R}$ to be an order preserving simplicial map which is an isomorphism on edges, thus the star $Y_i = \text{St}(v, X_i)$ is collapsed onto two consecutive edges $\overline{w_-w}$, $\overline{ww_+}$ in Y_{i-1}' , where $w = \pi_{i-1}(v)$, $w_- \prec w$, and $w \prec w_+$. Thus $\{Y_j\}$ is an admissible inverse system, but it need not satisfy (2) or (3) of Assumption 5.1.

Next, we enlarge $\{Y_j\}$ to a system $\{\hat{Y}_j\}$. We first attach, for every $k \geq i$, and every vertex $z \in (\pi_{i-1}^k)^{-1}(w_-)$, a directed ray γ_z which is directed isomorphic to $(-\infty, 0]$ with the usual subdivision and order. We then extend the projection maps so that if $\pi_k(z) = z'$ then $\gamma_z \subset Y_{j+1}$ is mapped direction-preserving isomorphically to a ray in X_j' starting at z' . Then similarly, we attach directed rays to vertices $z \in (\pi_{i-1}^k)^{-1}(w_+)$, and extend the projection maps.

Finally, we let $\{Z_j\}$ be the system obtained from $\{\hat{Y}_j\}$ by shifting indices by $(i - 1)$, in other words $Z_j = Y_{j-i+1}$.

Then $\{Z_j\}$ satisfies (2) and (3) of Assumption 5.1. For all $j \geq i$, we have compatible direction preserving simplicial embeddings $X_j \supset (\pi_i^j)^{-1}(\text{St}(v, X_i)) \rightarrow Z_{j-i+1}$. We will identify points with their image under this embedding. If $x, x' \in B(p, R)$ and $x = x_0, \dots, x_k = x'$ is a chain of points as in Lemma 2.3 which nearly realizes $\bar{d}_\infty^X(x, x')$, then the chain and the associated stars will project into $\text{St}(v, X_i)$; this implies that $\bar{d}_\infty^Z(x, x') \leq m^{(i-1)}\bar{d}_\infty^X(x, x')$. Similar reasoning gives $m^{(i-1)}\bar{d}_\infty^X(x, x') \leq \bar{d}_\infty^Z(x, x')$.

If $\gamma \subset B(p, R)$ is a monotone geodesic segment, then $\pi_i(\gamma)$ is a monotone geodesic segment in $\text{St}(v, X_i)$ with endpoints in $\text{St}(v, X_i)$, and so $\pi_i(\gamma) \subset \text{St}(v, X_i)$. Thus the embedding also preserves the partial order as claimed.

□

Fix $p \in X_\infty$. Then for every $n \in \mathbb{N}$, since $m^n < m^{(n+1)}$, Lemma 9.5 provides an inverse system $\{Z_j^n\}_{j \in \mathbb{Z}}$ and an embedding

$$\phi_n : (B(p, m^n), m^{-n-3}\bar{d}_\infty) \rightarrow Z_\infty^n.$$

Let $f_n : Z_\infty^n \rightarrow L_1$ be a 1-Lipschitz embedding satisfying the conclusion of Theorem 1.16, constructed in Section 9.1, and let $\psi_n : (B(p, m^n), \bar{d}_\infty) \rightarrow L_1$ be the composition $f_n \circ \phi_n$, rescaled by m^{n+3} . Next we use a standard argument with ultralimits, see [BL00]. Then the ultralimit

$$\omega\text{-lim } \psi_n : \omega\text{-lim}(B(p, m^n), \bar{d}_\infty) \rightarrow \omega\text{-lim } L_1$$

yields the desired 1-Lipschitz embedding, since X_∞ embeds canonically and isometrically in $\omega\text{-lim}(B(p, m^n), \bar{d}_\infty)$, and an ultralimit of a sequence of L_1 spaces is an L_1 space [Kak39].

9.3. Removing Assumption 5.1(3). Let $\{X_i\}$ be an admissible inverse system.

Lemma 9.6. *$\{X_i\}$ may be enlarged to an admissible inverse system $\{\hat{X}_i\}$ such that for all $i \in \mathbb{Z}$, \hat{X}_i is a union of monotone geodesics.*

Proof. We first enlarge X_i to \hat{X}_i as follows. For each $i \in \mathbb{Z}$, and each $v \in V_i$ which does not have a neighbor $w \in V_i$ with $w \prec v$ (respectively $v \prec w$), we attach a directed ray γ_v^- (respectively γ_v^+) which is directed isomorphic to $(-\infty, 0]$ (respectively $[0, \infty)$) with the usual subdivision and order. The resulting graphs \hat{X}_i have the property that every vertex $v \in \hat{X}_i'$ is the initial vertex of directed rays in both directions. Therefore we may extend the projection maps $\pi_i : X_{i+1} \rightarrow X_i$ by mapping $\gamma_v^\pm \subset X_{i+1}$ direction-preserving isomorphically to a ray starting at $\pi_i(v) \in X_i'$. The resulting inverse system is admissible. \square

If \bar{d}_∞^X and $\bar{d}_\infty^{\hat{X}}$ are the respective metrics, then for all $x, x' \in X_\infty \subset \hat{X}_\infty$, we clearly have $\bar{d}_\infty^{\hat{X}}(x, x') \leq \bar{d}_\infty^X(x, x')$. Note that if $x, x' \in X_\infty$ and $\{\pi_j(x), \pi_j(x')\}$ belong to the trimmed star of a vertex $v \in \hat{X}_j$, then in fact v is a vertex of X_j (since the trimmed star of a vertex in $\hat{X}_j \setminus X_j$ does not intersect X_j). Thus by Lemma 2.6 we have $\bar{d}_\infty^{\hat{X}}(x, x') \geq \frac{2m^2}{(m-2)} \bar{d}_\infty^X(x, x')$. Therefore if $f : \hat{X}_\infty \rightarrow L_1$ is the embedding given by Section 9.2, then the composition $X_\infty \hookrightarrow \hat{X}_\infty \xrightarrow{f} L_1$ satisfies the requirements of Theorem 1.16.

10. THE LAAKSO EXAMPLES FROM [Laa00] AND EXAMPLE 1.4

In [Laa00] Laakso constructed Ahlfors Q -regular metric spaces satisfying a Poincaré inequality for all $Q > 1$. In the section we will show that the simplest example from [Laa00] is isometric to Example 1.4.

10.1. Laakso's description. We will (more or less) follow Section 1 of [Laa00], in the special case that (in Laakso's notation) the Hausdorff dimension $Q = 1 + \frac{\log 2}{\log 3}$, $t = \frac{1}{3}$, and $K \subset [0, 1]$ is the middle third Cantor set.

Define $\phi_0 : K \rightarrow K$, $\phi_1 : K \rightarrow K$ by

$$\phi_0(x) = \frac{1}{3}x, \quad \phi_1(x) = \frac{2}{3} + \frac{1}{3}x.$$

Then ϕ_0 and ϕ_1 generate a semigroup of self-maps $K \rightarrow K$. Given a binary string $a = (a_1, \dots, a_k) \in \{0, 1\}^k$, we let $|a| = k$ denote its length. For every a , let $K_a \subset K$, be the image of K under the corresponding word in the the semigroup:

$$K_a = \phi_{a_1} \circ \dots \circ \phi_{a_k}(K).$$

Thus for every $k \in \mathbb{N}$ we have a decomposition of K into a disjoint union $K = \sqcup_{|a|=k} K_a$.

For each $k \in \mathbb{N}$, let $S_k \subset [0, 1]$ denote the set of $x \in [0, 1]$ with a finite ternary expansion $x = .m_1 \dots m_k$ where the last digit m_k is nonzero. In other words, if V_j is the set of vertices of the subdivision of $[0, 1]$ into intervals of length 3^{-j} for $j \geq 0$, then $S_k = V_k \setminus V_{k-1}$.

For each $k \in \mathbb{N}$ we define an equivalence relation \sim_k on $[0, 1] \times K$ as follows. For every $q \in S_k$, and every binary string $a = (a_1, \dots, a_k)$, we identify $\{q\} \times K_{(a_1, \dots, a_k, 0)}$ with $\{q\} \times K_{(a_1, \dots, a_k, 1)}$ by translation, or equivalently, for all $x \in K$, we identify $\phi_{a_1} \circ \dots \circ \phi_{a_k} \circ \phi_0(x)$ and $\phi_{a_1} \circ \dots \circ \phi_{a_k} \circ \phi_1(x)$.

Let \sim be the union of the equivalence relations $\{\sim_k\}_{k \in \mathbb{N}}$; this is an equivalence relation. We denote the collection of cosets $([0, 1] \times K) / \sim$ by F , equip it with the quotient topology, and let $\pi : [0, 1] \times K \rightarrow F$ be the canonical surjection. The distance function on F is defined by

$$d(x, x') = \inf \{ \mathcal{H}^1(\gamma) \mid \gamma \subset [0, 1] \times K, \pi(\gamma) \text{ contains a path from } x \text{ to } x' \},$$

where \mathcal{H}^1 denotes 1-dimensional Hausdorff measure.

10.2. Comparing F with Example 1.4. For every $k \in \mathbb{N}$ we will construct 1-Lipschitz maps $\iota_k : X_k \rightarrow F$, $f_k : F \rightarrow X_k$ such that $f_k \circ \iota_k = \text{id}_{X_k}$, such that the image of ι_k is $\text{const} \cdot 3^{-k}$ -dense in F . This implies that ι_k is an isometric embedding for all k , and is a $\text{const} \cdot 3^{-k}$ -Gromov-Hausdorff approximation. Therefore F is the Gromov-Hausdorff limit of the sequence $\{X_k\}$, and is isometric to (X_∞, d_∞) .

For every k , there is a quotient map $K \rightarrow \{0, 1\}^k$ which maps the subset $K_{(a_1, \dots, a_k)} \subset K$ to (a_1, \dots, a_k) . This induces quotient maps $K \times [0, 1] \rightarrow \{0, 1\}^k \times [0, 1]$, and $f_k : F \rightarrow X_k$, where X_k is the graph from Example 1.4. When X_k is equipped with the path metric described in the example, the map f_k is 1-Lipschitz, because any set $U \subset [0, 1] \times K$ with diameter $< 3^{-k}$ projects under the composition $[0, 1] \times K \rightarrow F \xrightarrow{f_k} X_k$ to a set $\bar{U} \subset X_k$ with $\text{diam}(\bar{U}) \leq \text{diam}(U)$.

For every k , there is an injective map $\{0, 1\}^k \rightarrow K$ which sends (a_1, \dots, a_k) to the smallest element of K_a , i.e. $\phi_{a_1} \circ \dots \circ \phi_{a_k}(0)$. This induces maps $[0, 1] \times \{0, 1\}^k \rightarrow [0, 1] \times K$ and $\iota_k : X_k \rightarrow F$. It follows from the definition of the metric on F that ι_k is 1-Lipschitz, since geodesics in X_k can be lifted piecewise isometrically to segments in $[0, 1] \times K$.

We have $f_k \circ \iota_k = \text{id}_{X_k}$. Therefore ι_k is an isometric embedding. Given $x \in [0, 1] \times K$, there exist $i \in \{0, \dots, 3^k\}$, $a \in \{0, 1\}^k$ such that $x \in W = [\frac{i-1}{3^k}, \frac{i}{3^k}] \times K_a$. Now W/\sim is a subset of F which intersects $\iota_k(X_k)$, and which has diameter $\leq 3^{-k} \text{diam}(F)$ due to the self-similarity of the equivalence relation, so ι_k is a $3^{-k} \text{diam}(F)$ -Gromov-Hausdorff approximation.

11. REALIZING METRIC SPACES AS INVERSE LIMITS: FURTHER GENERALIZATION

In this section we consider the realization problem in greater generality.

Let $f : Z \rightarrow Y$ be a 1-Lipschitz map between metric spaces. We assume that for all $r \in (0, \infty)$, if $U \subset Y$ and $\text{diam}(U) \leq r$, then the r -components of $f^{-1}(U)$ have diameter at most Cr .

Remark 11.1. Some variants of this assumption are essentially equivalent. Suppose $C_1, C_2, \bar{C}_1 \in (0, \infty)$. If for all $r \in (0, \infty)$ and every subset $U \subset Y$ with $\text{diam}(U) \leq r$, the $C_1 r$ -components of $f^{-1}(U)$ have diameter $\leq C_2 r$, it follows easily that the $\bar{C}_1 r$ -components of $f^{-1}(U)$ have diameter $\leq C_2 r \cdot \max(1, \frac{\bar{C}_1}{C_1})$.

11.1. Realization as an inverse limit of simplicial complexes. Fix $m \in (1, \infty)$ and $A \in (0, 1)$. For every $i \in \mathbb{Z}$, let \mathcal{U}_i be an open cover of Y such that for all $i \in \mathbb{Z}$:

- (1) The cover \mathcal{U}_{i+1} is a refinement of \mathcal{U}_i .
- (2) Every $U \in \mathcal{U}_i$ has diameter $\leq m^{-i}$.
- (3) For every $y \in Y$, the ball $B(y, Am^{-i})$ is contained in some $U \in \mathcal{U}_i$.

Next, for all $i \in \mathbb{Z}$ we let $f^{-1}(\mathcal{U}_i) = \{f^{-1}(U) \mid U \in \mathcal{U}_i\}$, and define $\hat{\mathcal{U}}_i$ to be the collection of pairs (\hat{U}, U) where $U \in \mathcal{U}_i$ and \hat{U} is an m^{-i} -component of $f^{-1}(U)$.

We obtain inverse systems of simplicial complexes $\{L_i = \text{Nerve}(\mathcal{U}_i)\}_{i \in \mathbb{Z}}$, and $\{K_i = \text{Nerve}(\hat{\mathcal{U}}_i)\}_{i \in \mathbb{Z}}$, where we view $\hat{\mathcal{U}}_i$ as an open cover of Z indexed by the elements of $\hat{\mathcal{U}}_i$. There are canonical simplicial maps $K_i \rightarrow L_i$ which send $(\hat{U}, U) \in \hat{\mathcal{U}}_i$ to $U \in \mathcal{U}_i$.

We may define a metric d_{K_∞} on the inverse limit K_∞ by taking the supremal metric on K_∞ such that for all $i \in \mathbb{Z}$ and every vertex $v \in K_i$, the inverse image of the closed star $\text{St}(v, K_i)$ under the projection $K_\infty \rightarrow K_i$ has diameter $\leq m^{-i}$. Let \bar{K}_∞ be the completion of (K_∞, d_{K_∞}) .

For every $z \in Z$ and $i \in \mathbb{Z}$, there is a canonical (possibly infinite dimensional) simplex σ_i in K_i corresponding to the collection of $U \in \hat{\mathcal{U}}_i$ which contain z . The inverse images $(\pi_i^\infty)^{-1}(\sigma_i) \subset \bar{K}_\infty$ form a nested sequence of subsets with diameter tending to zero, so they determine a unique point in the complete space \bar{K}_∞ . This defines a map $\phi : Z \rightarrow \bar{K}_\infty$.

Proposition 11.2. *ϕ is a bilipschitz homeomorphism.*

Proof. If $z, z' \in Z$ and $d(z, z') \leq Am^{-i}$, then $f(z), f(z') \in U$ for some $U \in \mathcal{U}_i$, and hence $z, z' \in \hat{U}$ for some m^{-i} -component $\hat{U} \in \hat{\mathcal{U}}_i$ of U . It follows that $d_{K_\infty}(\phi(z), \phi(z')) \leq m^{-i}$.

If $z, z' \in Z$ and $d_{K_\infty}(\phi(z), \phi(z')) \leq m^{-i}$, it follows from the definitions that $d(z, z') \lesssim m^{-i}$.

□

There is another metric \bar{d}_∞ on Z , namely the supremal metric with the property that every element of $\hat{\mathcal{U}}_i$ has diameter at most m^{-i} . Reasoning similar to the above shows that \bar{d}_∞ is comparable to d_Z .

11.2. Factoring f into “locally injective” maps. Let $\{\mathcal{U}_i\}_{i \in \mathbb{Z}}$ be a sequence of open covers as above.

For every $i \in \mathbb{Z}$, we may define a relation on Z by declaring that $z, z' \in Z$ are related if $f(z) = f(z')$ and $\{z, z'\} \subset \hat{U}$ for some $(\hat{U}, U) \in \hat{\mathcal{U}}_i$. We let \sim_i be the equivalence relation this generates. Note that \sim_{i+1} is a finer equivalence relation than \sim_i .

For every $i \in \mathbb{Z}$, we have a pseudo-distance d_i on Z defined by letting d_i be the supremal distance function $\leq d_Z$ such that $d_i(z, z') = 0$ whenever $z \sim_i z'$. Then $d_i \leq d_{i+1} \leq d_Z$, so we have a well-defined limiting distance function $d_\infty : Z \times Z \rightarrow [0, \infty)$. We let Z_i be the metric space obtained from (Z, d_i) by collapsing zero diameter subsets to points. We get an inverse system $\{Z_i\}_{i \in \mathbb{Z}}$ with 1-Lipschitz projection maps, and a compatible family of mappings $f_i : Z_i \rightarrow Y$ induced by f .

The map f_i is “injective at scale $\simeq m^{-i}$ ” in the following sense. If $z \in Z$, and $\bar{B} \subset Z_i$ is the image of the ball $B(z, Am^{-i})$ under the canonical projection map $Z \rightarrow Z_i$, then the restriction of f_i to \bar{B} is injective.

Proposition 11.3. *If $z, z' \in Z$ and $d_i(z, z') < m^{-i}$, then $d(z, z') \lesssim m^{-i}$. Consequently $d_\infty \simeq d_Z$.*

Proof. If $z_1, z_2 \in Z$ and $z_1 \sim_i z_2$, then z_1, z_2 belong to the same m^{-i} -component of $f^{-1}(\overline{B(f(z_1), 2m^{-i})})$, and hence $d(z_1, z_2) \leq 2Cm^{-i}$.

If $z, z' \in Z$ and $d_i(z, z') < m^{-i}$, then there are points $z = z_0, \dots, z_k = z'$ such if

$$J = \{j \in \{1, \dots, k\} \mid z_{j-1} \not\sim_i z_j\}$$

then

$$\sum_{j \in J} d(z_{j-1}, z_j) < m^{-i}.$$

Since f is 1-Lipschitz, it follows that $f(z_j) \in B(f(z), m^{-i})$ for all $j \in \{1, \dots, k\}$. Moreover, the z_j ’s lie in the same $2m^{-i}$ -component of $f^{-1}(B(f(z), m^{-i}))$, so $d(z, z') \leq 2Cm^{-i}$.

□

REFERENCES

- [Ass80] Patrice Assouad. Plongements isométriques dans L^1 : aspect analytique. In *Initiation Seminar on Analysis: G. Choquet-M. Rogalski-J. Saint-Raymond, 19th Year: 1979/1980*, volume 41 of *Publ. Math. Univ. Pierre et Marie Curie*, pages Exp. No. 14, 23. Univ. Paris VI, Paris, 1980.

- [BL00] Y. Benyamini and J. Lindenstrauss. *Geometric nonlinear functional analysis. Vol. 1*, volume 48 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2000.
- [Che99] J. Cheeger. Differentiability of Lipschitz functions on metric measure spaces. *Geom. Funct. Anal.*, 9(3):428–517, 1999.
- [CKa] J. Cheeger and B. Kleiner. in preparation.
- [CKb] J. Cheeger and B. Kleiner. Differentiating maps to L^1 and the geometry of BV functions. <http://arxiv.org/abs/math/0611954>.
- [CKc] J. Cheeger and B. Kleiner. Metric differentiation for PI spaces. In preparation.
- [CK06a] J. Cheeger and B. Kleiner. On the differentiability of Lipschitz maps from metric measure spaces into banach spaces. In *Inspired by S.S. Chern, A Memorial volume in honor of a great mathematician*, volume 11 of *Nankai tracts in Mathematics*, pages 129–152. World Scientific, Singapore, 2006.
- [CK06b] Jeff Cheeger and Bruce Kleiner. Generalized differential and bi-Lipschitz nonembedding in L^1 . *C. R. Math. Acad. Sci. Paris*, 343(5):297–301, 2006.
- [CK09] Jeff Cheeger and Bruce Kleiner. Differentiability of Lipschitz maps from metric measure spaces to Banach spaces with the Radon-Nikodým property. *Geom. Funct. Anal.*, 19(4):1017–1028, 2009.
- [CK10a] Jeff Cheeger and Bruce Kleiner. Differentiating maps into L^1 , and the geometry of BV functions. *Ann. of Math. (2)*, 171(2):1347–1385, 2010.
- [CK10b] Jeff Cheeger and Bruce Kleiner. Metric differentiation, monotonicity and maps to L^1 . *Invent. Math.*, 182(2):335–370, 2010.
- [CKN09] Jeff Cheeger, Bruce Kleiner, and Assaf Naor. A $(\log n)^{\Omega(1)}$ integrality gap for the sparsest cut SDP. In *2009 50th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2009)*, pages 555–564. IEEE Computer Soc., Los Alamitos, CA, 2009.
- [DCK72] D. Dacunha-Castelle and J. L. Krivine. Applications des ultraproducts à l'étude des espaces et des algèbres de Banach. *Studia Math.*, 41:315–334, 1972.
- [DL97] M. Deza and M. Laurent. *Geometry of cuts and metrics*, volume 15 of *Algorithms and Combinatorics*. Springer-Verlag, Berlin, 1997.
- [DU97] A. N. Dranishnikov and V. V. Uspenskij. Light maps and extensional dimension. *Topology Appl.*, 80(1-2):91–99, 1997.
- [Dyc74] R. Dyckhoff. Perfect light maps as inverse limits. *Quart. J. Math. Oxford Ser. (2)*, 25:441–449, 1974.
- [Eil34] S. Eilenberg. Sur les transformations continues d'espaces métriques compacts. *Fundamenta Mathematicae*, 1934.
- [Eng95] R. Engelking. *Theory of dimensions finite and infinite*, volume 10 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Lemgo, 1995.
- [HM82] S. Heinrich and P. Mankiewicz. Applications of ultrapowers to the uniform and Lipschitz classification of Banach spaces. *Studia Math.*, 73(3):225–251, 1982.
- [Kak39] S. Kakutani. Mean ergodic theorem in abstract (L) -spaces. *Proc. Imp. Acad., Tokyo*, 15:121–123, 1939.
- [Kir94] B. Kirchheim. Rectifiable metric spaces: local structure and regularity of the Hausdorff measure. *Proc. Amer. Math. Soc.*, 121(1):113–123, 1994.

- [Laa00] T. Laakso. Ahlfors Q -regular spaces with arbitrary $Q > 1$ admitting weak Poincaré inequality. *Geom. Funct. Anal.*, 10(1):111–123, 2000.
- [LP01] U. Lang and C. Plaut. Bilipschitz embeddings of metric spaces into space forms. *Geom. Dedicata*, 87(1-3):285–307, 2001.
- [Why34] G. T. Whyburn. Non-Alternating Transformations. *Amer. J. Math.*, 56(1-4):294–302, 1934.